

# Genetic Architecture of Tameness in a Rat Model of Animal Domestication

Frank W. Albert,<sup>\*1</sup> Örjan Carlborg,<sup>†</sup> Irina Plyusnina,<sup>‡</sup> Francois Besnier,<sup>†</sup> Daniela Hedwig,<sup>\*</sup>  
Susann Lautenschläger,<sup>\*</sup> Doreen Lorenz,<sup>\*</sup> Jenny McIntosh,<sup>\*</sup> Christof Neumann,<sup>\*</sup>  
Henning Richter,<sup>\*</sup> Claudia Zeising,<sup>\*</sup> Rimma Kozhemyakina,<sup>‡</sup> Olesya Shchepina,<sup>‡</sup>  
Jürgen Kratzsch,<sup>§</sup> Lyudmila Trut,<sup>‡</sup> Daniel Teupser,<sup>§</sup> Joachim Thiery,<sup>§</sup>  
Torsten Schöneberg,<sup>\*\*</sup> Leif Andersson,<sup>†,††</sup> and Svante Pääbo<sup>\*,††</sup>

<sup>\*</sup>Department of Evolutionary Genetics, Max Planck Institute for Evolutionary Anthropology, 04103 Leipzig, Germany, <sup>†</sup>Department of Animal Breeding and Genetics, Swedish University of Agricultural Sciences, 75007 Uppsala, Sweden, <sup>‡</sup>Institute of Cytology and Genetics, Siberian Branch of the Russian Academy of Sciences, 630090 Novosibirsk, Russia, <sup>§</sup>Institute of Laboratory Medicine, Clinical Chemistry and Molecular Diagnostics, University Hospital Leipzig, 04103 Leipzig, Germany, <sup>\*\*</sup>Institute of Biochemistry, Molecular Biochemistry, Medical Faculty, University of Leipzig, 04103 Leipzig, Germany, <sup>††</sup>Department of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden and <sup>†††</sup>Department of Genetics and Pathology, Rudbeck Laboratory, Uppsala University, 75123 Uppsala, Sweden

Manuscript received February 25, 2009

Accepted for publication April 3, 2009

## ABSTRACT

A common feature of domestic animals is tameness—*i.e.*, they tolerate and are unafraid of human presence and handling. To gain insight into the genetic basis of tameness and aggression, we studied an intercross between two lines of rats (*Rattus norvegicus*) selected over >60 generations for increased tameness and increased aggression against humans, respectively. We measured 45 traits, including tameness and aggression, anxiety-related traits, organ weights, and levels of serum components in >700 rats from an intercross population. Using 201 genetic markers, we identified two significant quantitative trait loci (QTL) for tameness. These loci overlap with QTL for adrenal gland weight and for anxiety-related traits and are part of a five-locus epistatic network influencing tameness. An additional QTL influences the occurrence of white coat spots, but shows no significant effect on tameness. The loci described here are important starting points for finding the genes that cause tameness in these rats and potentially in domestic animals in general.

**A**NIMAL domestication marked a turning point in human prehistory (DIAMOND 2002), and domestic animals have been the subject of research for many years (DARWIN 1868). Recently, genetic studies have shed light on when, where, and how often a range of animal species were domesticated (TROY *et al.* 2001; VILA *et al.* 2001; SAVOLAINEN *et al.* 2002; LARSON *et al.* 2005; DRISCOLL *et al.* 2007; ERIKSSON *et al.* 2008; NADERI *et al.* 2008). With the exception of coat color (*e.g.*, PIELBERG *et al.* 2008) and skin pigmentation (ERIKSSON *et al.* 2008), little is known about *what* occurred genetically during animal domestication. At what genes were allelic variants selected for by would-be practitioners of animal husbandry? Although domestic animals differ from each other in many ways, they all share the trait of tameness—*i.e.*, they tolerate and sometimes even seek human presence and handling. Almost nothing is currently known about the genetic basis of tameness.

In a series of studies initiated by D. K. Belyaev, researchers at the Institute for Cytology and Genetics in Novosibirsk (Russia) have subjected several mammalian species to a process of experimental domestication (TRUT 1999). These studies, some of them ongoing for several decades, involve selection for tame and aggressive behavior in lines of animals derived from wild populations. They include a fox population that has been “domesticated” to such an extent that the tame foxes are now similar to dogs in some respects (HARE *et al.* 2005). They also include a population of wild-caught rats (*Rattus norvegicus*) that was selected for either reduced or enhanced aggression toward humans over >60 generations (BELYAEV and BORODIN 1982). To select the animals, their response to an approaching human hand was observed, and the rats showing the least and the most aggressive behavior were allowed to mate within the two lines, respectively. The initial response to selection was rapid and then slowed, so that little change in behavior from generation to generation has been observed in the last 10–15 generations, although the selection regime has been continued to the present. Today, the “tame” rats are completely unafraid of humans, they tolerate handling

Supporting information is available online at <http://www.genetics.org/cgi/content/full/genetics.109.102186/DC1>.

<sup>1</sup>Corresponding author: Max Planck Institute for Evolutionary Anthropology, Evolutionary Genetics, Deutscher Platz 6, 04103 Leipzig, Germany. E-mail: falbert@eva.mpg.de

and being picked up, and they sometimes approach a human in a nonaggressive manner. By contrast, the “aggressive” rats ferociously attack or flee from an approaching human hand.

To study the genetic basis of tameness we have established populations of both rat lines in Leipzig. In their new environment, the rats maintained their behavioral differences in response to humans, and these differences were not influenced by postnatal maternal factors (ALBERT *et al.* 2008). In addition, the rat lines differ in a number of other behavioral, anatomical, and physiological traits, raising the question whether these traits are influenced by the same loci as tameness and aggression toward humans.

Many domestic animals display conspicuous coat color variations not found in their wild relatives. Prominent examples include the white color variants in dogs, pigs, cows, horses, and chickens. In laboratory rats, it has been proposed that “coat color genes” may account for many of the differences associated with domestication (KEELER and KING 1942). It is thus interesting that individuals with white spots appeared in both the tame foxes (TRUT 1999) and the tame rats (TRUT *et al.* 2000) at higher frequency than in the corresponding aggressive lines, although they were absent or rare in the founding fox and rat populations, and although they were not selected for. The rat populations studied here provide an excellent opportunity to examine whether tameness is influenced by the same loci as white coat spotting.

In this study, we crossed the two rat lines and bred >700 intercross animals. A broad set of behavioral, anatomical, and physiological traits was measured, and a genomewide set of genetic markers was used to identify genomic regions (quantitative trait loci, QTL) that influence tameness as well as other traits that differ between the lines, including white spots.

## MATERIALS AND METHODS

Additional materials and methods, as well as data files containing genotype and phenotype data collected for this study, can be found in the accompanying supporting information, File S1 and File S2.

**Animals:** The tame and the aggressive rat (*R. norvegicus*) lines derive from one population of wild-caught rats, which has been bidirectionally selected at every generation since 1972 at the Institute of Cytology and Genetics at the Siberian Branch of the Russian Academy of Sciences in Novosibirsk, Russia. About 30% of the animals from each generation were selected on the basis of the level of tameness and defensive aggression they displayed in response to humans on a five-point scale (NAUMENKO *et al.* 1989; PLYUSNINA and OSKINA 1997). Inbreeding was kept at a minimum by avoiding mating closely related individuals.

The pedigree described in this study was initiated from four tame and four aggressive individuals (2 females each; the “F minus one”, or “F<sub>-1</sub>”, generation) from the 64th generation of selection. The F<sub>-1</sub> animals did not have common parents and at most one common grandparent. They were mated

within line to yield 11 (5 tame and 6 aggressive rats, one male each) F<sub>0</sub> animals. These were crossed reciprocally, and six hybrid F<sub>1</sub> males were repeatedly mated to 37 F<sub>1</sub> females to produce 733 F<sub>2</sub> rats (383 females). A separate set of 47 F<sub>1</sub> animals (25 females) derived from different F<sub>0</sub> crosses was used for characterizing the F<sub>1</sub> generation in behavioral tests. Phenotypic data from F<sub>0</sub> animals discussed in this article are the same as presented in ALBERT *et al.* (2008). Animals were maintained under standard laboratory conditions, under an artificial 12-hr light cycle (lights off at 1:00 PM). The light cycle allowed behavioral testing during the dark phase, when rats are more active. Cages were equipped with sliding doors to allow for transfer without handling. During all caretaking procedures and experiments, animals from different lines and generations were treated identically. The study was approved by the regional government of Saxony (TVV Nr. 29/05).

**Behavioral testing:** F<sub>2</sub> rats were tested in a standardized series of behavioral paradigms. We measured the animals’ level of tameness/aggression with the “glove test,” by confronting them with an approaching human hand and attempting to handle them (see ALBERT *et al.* 2008 for details on the testing procedure). Beginning 2 weeks after the glove test, rats performed an open-field test, a light–dark test, and a startle response test, which provide various measures of exploratory and anxiety-related behavior. A total of 470 (64%) of the F<sub>2</sub> rats performed a second glove test trial. F<sub>1</sub> rats were tested once with the glove test at 12–14 weeks of age and then followed the testing schedule described in ALBERT *et al.* (2008). All tests were performed with minimal handling following procedures described in ALBERT *et al.* (2008). Glove test trials were videotaped and later analyzed by two independent observers (5% of trials only by one observer). Experimenters and observers were blind to the animals’ identity and to the further data processing. A set of 11 behaviors (*e.g.*, “attack” or “tolerate handling”) was scored (see ALBERT *et al.* 2008 for detailed descriptions of the behaviors), and each score was converted to a numeric measure (*e.g.*, “number of occurrences” or “duration in seconds”; Table 1).

**Blood and tissue sampling:** Dissections were performed on 733 F<sub>2</sub> (383 females) and 37 F<sub>1</sub> animals (16 females). Within 2 weeks after the last behavioral test, animals were killed between 2:00 and 6:00 PM. F<sub>2</sub> animals had been starved for 24 hr prior to dissection. Animals were weighed, anesthetized with CO<sub>2</sub>, and killed by cervical dislocation. Blood was collected rapidly after death by heart puncture and separated into serum and blood cells by centrifugation 30 min after sampling. Serum was stored at –20° and later transferred to –80° until analysis. Liver (small section, not weighed), spleen, kidney, adrenal gland, lung, and heart were removed rapidly, weighed, snap-frozen in liquid N<sub>2</sub>, and stored at –80°. From F<sub>1</sub> animals, kidney and spleen were weighed and stored.

**Serological phenotypes:** Serum was analyzed in 684 F<sub>2</sub> rats (357 females). Electrolytes, metabolites, immunological parameters, enzymes, and hormones were analyzed in serum according to the guidelines of the German Society of Clinical Chemistry and Laboratory Medicine. Measurements for all serum traits but corticosterone were performed using a Hitachi PPE-Modular analyzer (Roche Diagnostics, Mannheim, Germany). Corticosterone was measured using a commercial ELISA assay (IDS, Frankfurt, Germany).

**Statistical phenotype analyses:** We sought to control for possible confounding effects, such as observer in the glove test or an animal’s litter. We constructed mixed linear models of the phenotypes, estimated effect sizes using restricted maximum likelihood, and adjusted the phenotypes for the respective effects specified in Table S1. Sex and covariates were not adjusted; we instead included them as fixed effects into the QTL analyses. For glove test measures, we separately adjusted

observations from different trials and observers and then averaged the available observations from each rat. To summarize a rat's behavior in the glove test, we performed principal component analysis (PCA) on the individual glove test measures. We used only  $F_2$  animals in the PCA and calculated scores of  $F_0$  and  $F_1$  animals on the basis of the obtained regression coefficients.

Comparisons of non-glove test phenotypes between the  $F_0$ ,  $F_1$ , and  $F_2$  generations, as well as tests for sex differences (Wilcoxon's rank test), were performed on phenotypes adjusted for covariates. Correlations between phenotypes were calculated using Pearson's product-moment correlation, on phenotypes adjusted for all effects (including sex and covariates) listed in Table S1. All analyses were performed using the software R (R DEVELOPMENT CORE TEAM 2008) and the nlme (PINHEIRO *et al.* 2008) and lme4 (BATES 2007) packages.

**Genotyping:** Animals were genotyped with 152 microsatellite and 49 single-nucleotide polymorphism (SNP) markers. Markers were selected for maximum allele frequency differences between the outbred rat lines, as determined from preliminary genotyping of a panel of  $F_0$  animals. Preliminary genotyping of microsatellites was performed as described below, while SNPs were screened as described in SAAR *et al.* (2008). All markers used in the QTL mapping are listed in Table S2. DNA was isolated either from lung or from spleen tissue. Polymerase chain reaction (PCR) for microsatellite markers was performed using the M13-primer PCR system (SCHUELKE 2000) and analyzed on an ABI3730 DNA Analyzer (Applied Biosystems, Foster City, CA). Microsatellite genotypes were determined using the software GeneMapper Version 4.0 (Applied Biosystems), and all genotypes were manually double checked. SNPs were genotyped using TaqMan chemistry (Applied Biosystems). SNP genotypes were called automatically as part of the scanning process, and genotype plots were inspected visually.

**Pedigree construction:** Genotype data for individual markers were tested for Mendelian pedigree errors, using the program PedCheck (O'CONNELL and WEEKS 1998). Allele calls for inconsistent genotypes were rechecked manually and either they were corrected for obvious genotyping errors or the marker was excluded from further analysis if genotypes could not be determined unambiguously. We found that some rats yielded inconsistent genotypes for several markers, although the respective genotypes appeared to have been called correctly. We interpreted these individuals as having incorrect pedigree records. Initially, genotypes had been obtained from  $F_0$  and  $F_2$  animals. To clarify the pedigree structure, we typed all markers in all  $F_{-1}$  and  $F_1$  animals for which samples were available, as well as in an extended panel of  $F_0$  animals. Using the software "Cervus" 3.0 (KALINOWSKI *et al.* 2007), we determined the most likely parental pair for all genotyped  $F_0$ ,  $F_1$ , and  $F_2$  individuals, on the basis of a subset of 107 microsatellite markers with unambiguous genotype patterns. Rats for which the inferred parents differed from those in our records were either reassigned to the inferred parents or excluded from further analysis if no unique parental pair could be identified. The final pedigree used in the QTL analyses showed no Mendelian errors and comprised 8  $F_{-1}$  rats (all genotyped), 11  $F_0$  rats (8 genotyped), 43  $F_1$  rats (30 genotyped), and 706  $F_2$  rats (all genotyped).

**Linkage map construction:** We constructed a sex-averaged linkage map, using a version of the program crimap (GREEN *et al.* 1990) modified to handle large pedigrees by the University of California Davis, Veterinary Genetics Laboratory. On chromosome 6, we found the markers D6rat213 and D6rat68 to be inverted on our linkage map relative to their physical positions. On chromosome 13, the markers D13rat5 and D13rat64 were found to be inverted and to map very

closely (1.2 cM) to each other, despite a physical distance of 21 Mb. These cases may reflect chromosomal rearrangements in our wild-derived rats compared to the genome sequence of the inbred Brown Norway laboratory strain (GIBBS *et al.* 2004). We used our inferred linkage maps in further analyses (Table S2). We estimated information content at autosomal marker positions on the basis of the fraction of individuals whose alleles could be unambiguously traced to the  $F_{-1}$  generation.

**Single-QTL mapping:** A standard model of a phenotype  $y$  influenced by a single QTL can be written as

$$y = \beta_0 + FZ + \beta_1 a_j + \beta_2 d_j + \epsilon_j,$$

where  $\beta_0$  is the population mean,  $F$  is a matrix of regression coefficients for fixed effects and covariates (see Table S1 for the effects we included in the QTL models for various phenotypes),  $Z$  is an incidence matrix relating observations in  $F$  to individual observations,  $\beta_1$  and  $\beta_2$  are the additive and dominance effects at genomic position  $j$ ,  $a_j$  and  $d_j$  are indicator variables relating these genetic effects to individuals, and  $\epsilon_j$  is the residual error. We estimated the parameters  $\beta_0$ ,  $\beta_1$ , and  $\beta_2$  using a variation of the least-squares regression framework (HALEY and KNOTT 1992; HALEY *et al.* 1994). In this framework,  $F_2$  animals are grouped at a given genomic position according to whether they carry two, one, or zero alleles originated from the tame (allele "T") or aggressive (allele "A") line, forming the genotype classes TT, TA, and AA. Missing  $F_0$  genotypes can lead to a loss of power because some alleles in  $F_2$  animals might not be reliably traced to parental lines. This limitation can be overcome by including the genotyped parents of  $F_0$  animals (the  $F_{-1}$ ) in the analysis and by tracing alleles back to them. Hence, we inferred missing genotypes of  $F_0$  and  $F_1$  animals on the basis of their ancestors' and offspring genotypes (File S1). Next, we calculated, in steps of 1 cM throughout the genome, the probability of an  $F_2$  animal belonging to the TT, TA, or AA genotype classes on the basis of genotypes of flanking markers, using methods described in PONG-WONG *et al.* (2001) and BESNIER and CARLBORG (2007). The genotype probabilities were used to compute the indicator variables  $a_j$  and  $d_j$  (HALEY *et al.* 1994). Finally, we fitted the parameters  $\beta_0$ ,  $\beta_1$ , and  $\beta_2$  using least-squares regression. High  $F$ -values obtained from the regression point to the presence of a putative QTL at the respective location. The difference in mean between the two homozygous classes TT and AA corresponds to twice the additive effect  $\beta_1$  of the QTL, while the deviation of the heterozygous TA class from the mean of the TT and AA classes measures the degree of dominance  $\beta_2$ . We searched for QTL by following a forward selection procedure (Figure S1). After the initial scan, we included the effect and position of the most significant QTL that reached genomewide significance as a fixed effect in the model. The scan was then repeated, and QTL were added until no additional significant QTL were identified. For each phenotype, we performed 1000 permutations of phenotypes with regard to genotypes to determine the  $F$ -value threshold that corresponds to a genomewide significance level of  $P = 0.05$  (CHURCHILL and DOERGE 1994). We express variance components attributable to QTL as a fraction of the residual phenotypic variance, *i.e.*, the variance in phenotype after fixed effects and covariates have been factored out.

We analyzed the X chromosome using the software QxPak (PEREZ-ENCISO and MISZTAL 2004) (see File S1 for details). Since the permutation-based significance thresholds derived for the autosomes cannot be directly applied to the X chromosome, we assumed QTL with a nominal  $P$ -value  $< 0.001$  (0.005) to be significant (suggestive) at a genomewide level, as suggested in the QxPak manual. We tested for linkage to the Y chromosome, using ANOVA as described in File S1.



**Mapping of epistatic QTL:** We searched for epistatic QTL using an extension of the least-squares regression model for single QTL, following a search strategy described and applied earlier (CARLBORG and ANDERSSON 2002; CARLBORG *et al.* 2003, 2004, 2006; WRIGHT *et al.* 2006). Here, we first describe the regression model underlying our epistatic analyses and then go through the steps of the search strategy. A schematic representation of the approach is shown in Figure S1.

The standard extension of the model for a single QTL to incorporate two epistatic QTL is

$$y = \beta_0 + FZ + \beta_{1_j} a_j + \beta_{2_j} d_j + \beta_{3_k} a_k + \beta_{4_k} d_k + \beta_{5_{jk}} aa_{jk} + \beta_{6_{jk}} ad_{jk} + \beta_{7_{jk}} da_{jk} + \beta_{8_{jk}} dd_{jk} + \varepsilon_{jk}.$$

The additional parameters are the additive ( $\beta_{3_k}$ ) and dominance effects ( $\beta_{4_k}$ ) of the second QTL at position  $k$ , the epistatic effects between the two loci ( $\beta_{5_{jk}}-\beta_{8_{jk}}$ ), and the corresponding indicator variables. The indicator variables for the interaction terms ( $aa_{jk}$ ,  $ad_{jk}$ ,  $da_{jk}$ ,  $dd_{jk}$ ) were computed by multiplying the respective additive and dominance indicator variables for the single QTL (HALEY and KNOTT 1992). We estimated the effects  $\beta_0-\beta_8$  at a given pair of loci using least-squares regression.

The search strategy to find epistatic pairs of QTL involves three steps (see Figure S1 and CARLBORG *et al.* 2004 for further details). First, we searched for single QTL as described above. Second, we performed a genomewide search for putative epistatic pairs of loci. Third, each pair is tested for the existence of epistasis.

For each pair proposed by the pairwise search in step two, we assigned significance in one of three ways. If both loci were significant by themselves in the single-QTL analysis, no further testing is necessary, and the pair is declared significant. If one locus in the putative pair was a significant single QTL, we need to test only whether the second locus is significant (*i.e.*, whether its inclusion in the model already containing the first locus improves model fit). To derive the corresponding threshold, we created, for each single QTL, 1000 randomized data sets by permuting only the indicator variables of the second QTL ( $a_k$ ,  $d_k$ ), as well as those of the interaction effects ( $aa_{jk}$ ,  $ad_{jk}$ ,  $da_{jk}$ ,  $dd_{jk}$ ), while the first QTL was kept in the model as a fixed effect (permutation test “Type I” in Figure S1). In each permuted data set, we searched for the best fitting pair including the known QTL and a second QTL. We then compared the model fit obtained from the putative pair with the model fits obtained from the permutations. Finally, if neither QTL in the putative pair was significant as a single QTL, we need to test whether the joint inclusion of both loci improves the model fit significantly. We performed 1000 phenotype/genotype permutations and searched each randomized data set for its best fitting pair (permutation test “Type II” in Figure S1). For increased efficiency, this was done using a genetic search algorithm (CARLBORG *et al.* 2000). Significance of the putative pair was assigned by comparing its model fit to the fits obtained from the permutations. Throughout, we used a cutoff of  $P < 0.05$  for “significant” pairs of QTL.

So far, we have detected pairs of QTL, but not yet tested whether there is significant epistasis between the members of a given pair. To do this, we generated 1000 randomized data sets for each pair by permuting only the interaction indicator variables ( $aa_{jk}$ ,  $ad_{jk}$ ,  $da_{jk}$ ,  $dd_{jk}$ ), while keeping the additive and dominance effects of the two loci in the pair constant (CARLBORG and ANDERSSON 2002) (permutation test “Type III” in Figure S1). Epistasis is assumed if the putative epistatic pair is in the top 5% of model fits obtained from the permutations.

To construct the network influencing tameness, we consider loci (single or as part of a pair) with overlapping confidence

intervals to be the same locus. We show these loci as circles in Figure 5, with lines between them indicating significant epistatic interactions for that given pair. We did not fit a model incorporating interactions between more than two loci. To visualize the directions of the epistatic interactions (Figure 5, B–F), we grouped  $F_2$  animals according to their genotypes at the respective loci. For each of the resulting nine two-locus genotype groups, we calculated the mean and the standard error of the mean of the respective animals’ level of tameness.

Because some loci in the tameness network are part of more than one epistatic interaction, we cannot calculate the residual phenotypic variance explained by the whole network by simply adding up the variances explained by the respective pairs (Table 4). Instead, we used the NOIA model of genetic effects (ALVAREZ-CASTRO and CARLBORG 2007) as implemented in LE ROUZIC and ALVAREZ-CASTRO (2008), with analyses restricted to at most pairwise interactions. NOIA is specifically designed to estimate parameters, including genetic variances, in multi-locus networks (ALVAREZ-CASTRO and CARLBORG 2007).

## RESULTS

**A cross between tame and aggressive rats:** To create an intercross between the tame and aggressive rats, we mated one tame and one aggressive male to 5 aggressive and 4 tame females, respectively. In the resulting  $F_1$  generation, we repeatedly mated six males with a total of 37 females to produce an  $F_2$  population of 733 animals (362 females). Details of the mating scheme are described in MATERIALS AND METHODS.

**Analyses of phenotypes:** We recorded a total of 45 phenotypic traits in the  $F_2$  animals, including measures from four behavioral tests, anatomical parameters, and serum levels of hormones, enzymes, and other serum components (Table 1). To measure the level of tameness and aggression, we used a paradigm that closely mimics the test used to select the two rat lines over the past 36 years. In this “glove test,” a gloved human hand approaches a rat in an experimental cage and attempts to touch it and to pick it up (Figure 1A). Various aspects of the rat’s behavior are recorded (ALBERT *et al.* 2008). When testing  $F_1$  animals in this test, we found that the extreme levels of tameness and aggression observed in the original  $F_0$  lines were absent (Figure S2). By contrast,  $F_2$  animals displayed the full range of behaviors found in the original lines (Figure S2). A few  $F_2$  animals even exceeded the levels of tameness and aggression observed in the tame and aggressive lines.

A PCA of the behaviors recorded in the glove tests of the  $F_2$  animals confirms these observations. The first principal component (PC1) corresponds to behaviors such as attacks, screaming, and (with inverse loading) the toleration of touch or handling (Table 2). PC1 explains 26% of the variance in behavior of  $F_2$  rats. Of the PCs explaining  $>10\%$  of the variance, PC1 most clearly separates the tame from the aggressive  $F_0$  animals (Wilcoxon’s rank test: PC1,  $P < 10^{-15}$ ; PC2,  $P = 0.003$ ; PC3,  $P = 0.07$ ), although these animals were not included in the PCA (see MATERIALS AND METHODS).

**TABLE 1**  
**Traits measured in F<sub>2</sub> animals**

Trait	Unit	No. F <sub>2</sub>	Higher trait value	Comments
Glove test		700		
Approach	Count		Tame	No. of occurrences
Attack	Count		Aggressive	No. of occurrences
Escape	Count		Aggressive	No. of occurrences
Flight	Count		Aggressive	No. of occurrences
Move and leave	Count		Tame	No. of occurrences
Squeak	Count		Aggressive	No. of occurrences
Boxing	sec		Aggressive	Duration
Freeze	sec		—	Duration
Scream	sec		Aggressive	Duration
Tolerate handling	sec		Tame	Duration
Tolerate touch	sec		Tame	Duration
Open field test		689		
Time spent in center	%		Tame	
Time spent in corner	%		Aggressive	
Time spent moving	%		Tame	
Time spent rearing	sec		Tame	
Locomotion speed	cm/sec		Tame	
Fecal boli	Count		—	
Light–dark test		690		
Time spent in light compartment	%		Tame	
Time spent moving	%		—	
Time spent rearing	sec		Tame	
Locomotion speed in light compartment	cm/sec		—	
Fecal boli	Count		—	
Startle response test		700		
Startle response	g		Aggressive	Mean startle response across 10 trials
Anatomy				
Body weight	g	700	—	
Adrenal gland weight	g	664	Aggressive	
Heart weight	g	669	—	
Kidney weight	g	669	Tame	
Lung weight	g	668	—	
Spleen weight	g	668	Tame	
Testis weight	g	321	—	
White coat spotting	Yes/no	573	Tame	
Serum traits		638		
Hormones				
Corticosterone	ng/ml		Aggressive	
fT3	pmol/liter		—	
fT4	pmol/liter		Aggressive	
Enzymes				
ALAT	μkat/liter		Aggressive	
AP	μkat/liter		Aggressive	
ASAT	μkat/liter		Aggressive	
Substrates				
Creatinine	μmol/liter		Aggressive	
Glucose	μmol/liter		Aggressive	
Urea	mmol/liter		Tame	
Serum protein				
Total protein	g/liter		Aggressive	
Albumin	g/liter		Tame	
Electrolytes				
Ca <sup>2+</sup>	mmol/liter		Tame	
Cl <sup>-</sup>	mmol/liter		Aggressive	
Fe <sup>2+/3+</sup>	μmol/liter		Tame	

ALAT, alanine aminotransferase; AP, alkaline phosphatase; ASAT, aspartate aminotransferase; fT3, free triiodothyronine; fT4, free thyroxine.




FIGURE 1.—Behavior in the glove test. (A) The glove test measures the level of tameness or aggression toward an approaching hand. (B) PCA scores derived from glove test behaviors of tame  $F_0$  (blue), aggressive  $F_0$  (red),  $F_1$  (purple), and  $F_2$  (black) animals. Circles, females; squares, males.

Of the  $F_1$  rats, 94% (44/47) had PC1 scores between the medians of the tame and the aggressive rats. Of the  $F_2$  rats, 79% (551/700), fell into this intermediate range, while 6% (43) of the  $F_2$  rats had more tame and 16% (109) had more aggressive PC1 scores than defined by the respective  $F_0$  medians (Figure 1B). This indicates that there is substantial variation in tameness in the  $F_2$  rats and suggests that PC1 is a useful measure of this variation. The glove tests were repeated in 470  $F_2$  rats (MATERIALS AND METHODS). The correlation between the PC1 scores obtained for the two trials was 0.44 (Pearson's  $r$ ;  $P < 10^{-15}$ ).

We also performed an open-field test, a light–dark test, and a startle response test, which measure traits related to anxiety and fear as well as general activity (Table 1). The values of all these traits in  $F_2$  animals overlapped substantially with those in  $F_0$  animals (Figure S3). This was also true for body weight, for the weight of six organs, and for 8 of 14 serum traits (Figure S4 and Figure S5). By contrast, >75% of the  $F_2$  rats had higher (corticosterone, creatinine, glucose, chloride) or lower (alanine aminotransferase, ALAT; alkaline phosphatase, AP) values in these measures than >75% of the tame and aggressive rats (Figure S5).

Sex differences were apparent for many traits in the  $F_2$  animals (Table S3). When males and females were

TABLE 2

Principal component analysis of behavior of  $F_2$  animals in the glove test

Measure	PC1	PC2	PC3
Attacks (count)	−0.43		
Boxing posture (duration)	−0.36		
Escapes (count)	−0.32		
Flights (count)	−0.39		
Screaming (duration)	−0.38		
Freezing (duration)		0.46	−0.36
Move and leave (count)		−0.42	−0.61
Squeaks (count)		−0.35	0.31
Approaches (count)		−0.43	
Tolerate handling (duration)		0.41	0.50
Tolerate touch (duration)	0.46		
% variance	26	15	11

The loadings shown for the respective principal components (PCs) indicate the degree to which a trait contributes to the respective PC. Only PCs that explain  $\geq 10\%$  of the variance and loadings with absolute values  $\geq 0.3$  are shown.

considered separately, most phenotypes were approximately normally distributed in the  $F_2$  generation (Figure S6, Figure S7, Figure S8, and Figure S9). By contrast, raw glove test measures had highly skewed distributions, with prominent peaks at zero counts/durations.

**Correlations between phenotypes:** Earlier work revealed a multitude of phenotypic differences between the tame and the aggressive rats, including behavioral, anatomical, hormonal, and neurochemical differences (NAUMENKO *et al.* 1989; PLYUSNINA and OSKINA 1997; POPOVA *et al.* 2005; ALBERT *et al.* 2008). If these differences are caused by the same genetic loci, they should be correlated in the  $F_2$  animals. We did observe significant correlations among parameters recorded in the same test. For example, correlations between different measures in the glove test are reflected in their contributions to PC1 (Table 2). In contrast, parameters from different tests were generally not, or only weakly, correlated (Figure 2, Table S4). Notably, two measures of “anxiety” (the percentage of time spent in the light compartment of the light–dark test and the percentage of time spent in the center of the open field) were not correlated ( $r = 0.03$ ,  $P = 0.48$ ). There were significant but weak correlations between tameness and aspects of behavior in the open-field and light–dark tests. High levels of tameness were also correlated with low corticosterone levels ( $r = -0.08$ ,  $P = 0.04$ ), but not with the weight of the adrenal glands ( $r = -0.06$ ,  $P = 0.10$ ).

**QTL for tameness and associated traits:** We typed the animals in the pedigree for 201 genetic markers (152 microsatellites and 49 single-nucleotide polymorphisms) that were selected to be polymorphic between the parental strains and to provide coverage of most of the genome (MATERIALS AND METHODS).






FIGURE 2.—Correlations between phenotypes. Positive correlation coefficients are shown in red and negative ones in blue. Red boxes mark correlations within the same test or group of traits.

The phenotypic and genetic data were used to identify QTL for the traits measured in the  $F_2$  animals. A total of 23 significant and 125 suggestive autosomal QTL, and one significant QTL on the X chromosome, were identified when analyzing both sexes together (Table 3 and Table S5 and Table S6). All but two serum traits (FT3 and calcium levels) showed at least suggestive QTL. In what follows, we focus on QTL for tameness and overlapping QTL, as well as one QTL for coat color.

Two significant QTL for tameness (measured as PC1) were identified (Figure 3A). The strongest of these (termed “*Tame-1*”) is located at 58 cM on chromosome 1. The difference in tameness between homozygous genotypes at *Tame-1* corresponds to  $\sim 20\%$  of the difference between the tame and the aggressive line (1.2 of 6.4 units of PC1). *Tame-1* explains 5.1% of the phenotypic variance in tameness. The second locus (“*Tame-2*”) is located at 78 cM on chromosome 8. Both the tameness

difference between homozygous genotypes ( $\sim 10\%$  of the line difference) and the portion of residual phenotypic variance in tameness it explains (2.3%, are about half of those of *Tame-1*.

The region encompassed by *Tame-1* also contains significant QTL for rearing in the open field and for adrenal gland weight, as well as a suggestive QTL for the time spent moving in the open field (Figure 3B). The effects of these QTL are in the expected direction—*i.e.*, alleles from the tame line influence the phenotype in the direction expected from the comparison between tame and aggressive animals (*e.g.*, causing higher tameness scores and lower adrenal gland weight). However, *Tame-1* also overlaps with a significant QTL that influences spleen weight. This QTL is transgressive—*i.e.*, the alleles from the tame line reduce spleen weight although the rats from the tame line have 30% heavier spleens on average (ALBERT *et al.* 2008).

**TABLE 3**  
**Autosomal QTL identified at genomewide significance**

Trait	Chr <sup>a</sup>	Peak <sup>b</sup>	1 LOD C.I. <sup>b,c</sup>	F	Additive effect	Dominance	% variance <sup>d</sup>	Females <sup>e</sup>	Males <sup>e</sup>
Glove test									
Tameness (PC1)	1	58	54–64	19	0.60	0.43	5.1	Significant	Suggestive
Tameness (PC1)	8	78	68–85	9	0.37	0.43	2.3	—	Suggestive
Attack	1	60	51–73	14	–0.55	–0.48	3.8	Suggestive	—
Flight	1	48	42–64	11	–0.36	–0.08	3.2	Significant	—
Screaming	1	69	64–80	10	–0.80 sec	–0.97 sec	3.8	—	Significant
Tolerate touch	1	53	50–60	17	1.6 sec	1.0 sec	4.6	Significant	—
Other behavioral tests									
Time in corner (OF)	8	90	85–96	8	–0.05%	–0.02%	2.4	Suggestive	Suggestive
Rearing (OF)	1	72	66–83	10	3.1 sec	–0.4 sec	2.7	Suggestive	—
Startle response	10	68	57–78	11	–66 g	–7.5 g	3.0	—	Suggestive
Anatomy									
Adrenal gland	1	55	52–60	42	–3.8 mg	–0.6 mg	11.3	Significant	Significant
Adrenal gland	20	34	21–42	10	–0.6 mg	–2.0 mg	2.5	Suggestive	Suggestive
Body weight	7	30	14–46	10	11 g	5.3 g	2.9	Suggestive	Suggestive
Body weight	8	96	92–97	9	10 g	1.4 g	2.3	Suggestive	—
Heart	3	83	74–90	14	39 mg	64 mg	3.9	Suggestive	Suggestive
Spleen	10	68	58–72	16	25 mg	1.1 mg	4.6	Suggestive	Suggestive
Spleen	1	85	60–93	11	–20 mg	7.8 mg	3.0	Suggestive	Suggestive
Spleen	5	114	107–114	9	–17 mg	–10 mg	2.4	—	—
White spotting	14	26	21–33	22	0.20	–0.07	7.1	Significant	Significant
Serum traits									
Albumin	6	87	75–93	10	–0.6 g/liter	–0.3 g/liter	3.1	—	Suggestive
AP	11	22	4–35	9	0.07 $\mu$ kat/liter	–0.02 $\mu$ kat/liter	2.8	Suggestive	—
AP	5	114	98–114	8	–0.060 $\mu$ kat/liter	0.008 $\mu$ kat/liter	2.5	—	Significant
Creatinine	2	90	82–108	11	1.3 $\mu$ mol/liter	0.3 $\mu$ kat/liter	3.3	—	Suggestive

OF, open field test.

<sup>a</sup>Chromosome.

<sup>b</sup>Centimorgans.

<sup>c</sup>Confidence interval.

<sup>d</sup>Residual phenotypic variance explained after accounting for fixed effects.

<sup>e</sup>Significance level when analyzing females/males separately.

In the proximity of *Tame-2* several other QTL are found, two of which are significant. At these QTL, the alleles from the tame line increase body weight and decrease the time an animal spent in the corners of the open field, respectively. These effects are in the expected direction. Among suggestive QTL, those for time spent in the center, time spent moving and the number of rears in the open field, kidney weight, and serum aspartate aminotransferase (ASAT) all have effects in the expected direction. There were also two transgressive suggestive QTL, where the tame allele increases adrenal gland weight and the magnitude of the startle response, respectively.

We found white coat spotting to be linked to a significant QTL on chromosome 14, but no aspect of tameness mapped to this region (Figure 4). White coat spotting did not show association to *Tame-1*, *Tame-2* (Figure S10), or any other QTL for behavior during the glove test. Further, the tameness levels of individuals carrying white ventral spots did not differ significantly from those without them (*t*-test,  $P = 0.17$ , Figure 4A).

To assess whether the QTL we identified might be specific to one sex, we analyzed all traits using only

female or male F<sub>2</sub> animals, respectively (Table 3 and Table S5 and Table S7). For tameness, *Tame-1* reached genomewide significance in females and chromosome-wide significance in males, where the *F*-value (8.6) was close to the genomewide significance threshold (8.8). *Tame-2* reached chromosome-wide significance only in males. The QTL for adrenal gland weight on chromosome 1 and for white spotting on chromosome 14 were significant in both sexes. Several QTL were suggestive in both sexes (*e.g.*, spleen weight on chromosomes 1 and 10), whereas for several others we found significant or suggestive linkage only in one sex. All individual behaviors in the glove test, but not PC1, fall in the latter category. For example, at *Tame-1*, only males showed a significant QTL for screaming, whereas only females had significant QTL for flights and toleration of touch. Toleration of touch yielded one additional significant sex-specific QTL in females and males, respectively (Table S5).

**Epistatic interactions:** Epistatic interactions can have large effects on phenotypic traits. Hence, we searched the genome for interacting pairs of loci for all traits described in this study. Fifteen epistatic pairs affecting






FIGURE 3.—QTL for tameness and aggression. (A) Evidence for linkage to tameness across the genome. High  $F$ -values indicate the presence of a QTL. The dashed horizontal line represents the genomewide significance threshold. Solid horizontal bars are 2-LOD drop confidence intervals for QTL position. Chromosome boundaries are indicated by upward tick marks on the  $x$ -axis. (B) Several traits map to the two QTL for tameness. Only significant QTL are shown. Black and red upward tick marks are microsatellite and SNP marker positions, respectively.

9 traits reached genomewide significance ( $P < 0.05$ ), exceeding the random expectation of  $<3$  pairs for 48 analyzed traits. Most pairs were found for tameness, forming an interconnected network of 5 loci (Figure 5A). The network explained 14% of the residual phenotypic variance, compared to 7.4% explained by *Tame-1* and *Tame-2* individually. It is discussed below, while epistatic pairs for the remaining traits are given in Table 4.

The tameness network comprises five pairwise interactions between five loci (Figure 5A). Two loci in the network had significant individual effects (*Tame-1* and *Tame-2*). When considering these loci simultaneously, the tame allele (T) at locus *Tame-1* increases tameness regardless of the *Tame-2* genotype (Figure 5D). The effect is, however, strongest when *Tame-2* is homozygous (AA) for the allele from the aggressive line (A). The effect of the tame allele at *Tame-2* is smallest when *Tame-1* is homozygous for the allele from the tame line (TT), where the difference between the three *Tame-2* genotypes (AA, AT, and TT) is not significant.

Both *Tame-1* and *Tame-2* interact with a locus on chromosome 19 (denoted *Tame-3*) that reached only suggestive significance in the scan for single QTL. The tame allele at *Tame-1* has a significant effect on tameness only in a heterozygous (AT) background of *Tame-3*

(Figure 5B). There is no additive effect of *Tame-3* on tameness, but strong overdominance for aggression in the *Tame-1* AA genetic background and a strong overdominance for tameness in the *Tame-1* TT genetic background. This interesting shift in the direction of the dominance effect deserves further investigation. When considering *Tame-3* and *Tame-2* together, the only genotype with a deviating phenotype is the AAAA double homozygote that significantly increases aggression (Figure 5C). In addition, *Tame-1* and *Tame-2* interact significantly with one additional locus each. The effect of the tame allele at *Tame-1* is strongest when a locus on chromosome 4 (denoted *Tame-4*) is TT, and the tame allele at *Tame-4* has a significant effect on tameness only in the AA *Tame-1* genotype (Figure 5E). In this background, it is transgressive, increasing aggression. The *Tame-2* genotype also has a major effect on a locus on chromosome 6 (denoted *Tame-5*), in that *Tame-5* affects tameness only when *Tame-2* is AA (Figure 5F).

## DISCUSSION

**A polygenic basis for tameness:** To uncover the genetic basis for tameness, we analyzed  $>700$   $F_2$  animals produced by cross-breeding two strains of rats that differ drastically in their response to humans. Tameness levels,




FIGURE 4.—White coat spotting and tameness. (A) Tameness level of  $F_2$  animals with ( $n = 190$ ) and without ( $n = 393$ ) white ventral coat spots. (B) A QTL for spotting (black line) does not show linkage to tameness (red line). The vertical dashed line indicates the location of the *Kit* gene. Black and red upward tick marks are microsatellite and SNP positions, respectively.

as measured by the glove test in the  $F_1$  and  $F_2$  rats, approximate a normal distribution with the mean centered between those of the parental lines (Figure 1), suggesting a polygenic basis for tameness and aggression in the rats (LYNCH and WALSH 1998).

The linkage study confirms this. The largest QTL, *Tame-1*, explains 5.1% of the residual phenotypic variance, while the remaining loci each explain smaller fractions when considered individually. These estimates are in line with the generally small effect sizes reported for QTL for other rodent behaviors (FLINT 2003). Across 45 measures and three principal component scores, our study identified 23 significant and 125

suggestive QTL. We note that this greatly exceeds the number of QTL expected to be observed by chance. For 48 traits, we expect  $<3$  significant QTL at a genomewide significance level of 5% and 48 suggestive QTL with a significance level of 5% at each of 20 chromosomes. Given the sample size of  $>700$   $F_2$  animals, we consider it unlikely that other unidentified loci with large individual effects exist in these lines.

**Overlap of QTL for tameness-associated traits:** Previous studies have identified a multitude of phenotypic differences between the tame and the aggressive lines of rats (NAUMENKO *et al.* 1989; PLYUSNINA and OSKINA 1997; POPOVA *et al.* 2005; ALBERT *et al.* 2008). However, the behavioral response to humans was the only criterion used during selection. Are the loci influencing the nonselected traits the same as those contributing to the difference in tameness?

If phenotypic traits are influenced by the same loci one would expect them to show some degree of correlation. It is thus noteworthy that correlations between the traits we measured in the  $F_2$  population were weak at best, often failing to reach significance in spite of the fact that hundreds of animals were analyzed and that some of the uncorrelated traits were markedly different between the parental strains (ALBERT *et al.* 2008). However, the power to detect a correlation caused by shared loci may be limited given that the effect sizes associated with the alleles are small and perhaps obscured by nongenetic influences.

In the QTL analyses, a number of traits mapped to the same regions. This is especially obvious on chromosome 1 where weight of the adrenal gland maps to a region overlapping *Tame-1* with virtually identical confidence intervals (Figure 3B). It thus seems plausible that alleles of a single gene with pleiotropic effects underlie both *Tame-1* and adrenal gland size variation. Alternatively, a causal relationship might exist between tameness and adrenal gland size. For example, sudden increases of plasma corticosterone, which is produced by the adrenal cortex, promote aggressive behavior in rats (KRUK *et al.* 2004), while chronically high levels of glucocorticoids seem to inhibit aggressive behavior in several vertebrate species (SUMMERS *et al.* 2005). However, *postmortem* corticosterone levels did not map to any locus linked to tameness or aggression. It is thus equally possible that other hormonal activities of the adrenal glands are directly or indirectly linked to the alleles that affect tameness in the rats. The identification of the gene or genes underlying *Tame-1* as well as adrenal gland size variation will eventually clarify this.

*Tame-1* was obtained by mapping a principal component score summarizing, among other traits, attacks, screaming, flights, and the toleration of touch (Table 2). The fact that, at *Tame-1*, these individual behaviors yielded QTL apparently specific to one sex (Table 3) may suggest that the causative alleles underlying *Tame-1* influence the two




FIGURE 5.—An epistatic network for tameness. (A) Overview of QTL (circles) and epistatic interactions (lines). Only QTL in pairs with significant epistatic interactions are shown. Bold (nonbold) solid circles: the QTL was significant (suggestive) in the scan for single loci. Dashed circles: the QTL was significant only as part of an epistatic pair. Numbers in circles indicate QTL chromosome and position (centimorgans). (B–F) Phenotypes for two-locus genotypes. Circles indicate the mean phenotype for the respective two-locus genotypes; error bars show the standard error of the mean. The strength of the QTL effect at the first locus in the pair is indicated by the slope of the line connecting the homozygous genotypes [both alleles from the tame line (TT) vs. both alleles from the aggressive line (AA)]. An effect at the second locus in the pair is indicated by nonoverlapping allele effects at a given genotype of the first locus.

sexes differently. However, since the QTL for the principal component at *Tame-1* is found in both sexes, it may be the case that the absence of signals for the individual traits reflects lower power due to using half the number of individuals. In addition, individual traits are likely to have less power than the principal component they contribute to. Further work is warranted to clarify whether *Tame-1* and other loci truly act in a sex-specific manner.

To our knowledge, this study is the first genetic mapping of tameness and defensive aggression against humans in any species. However, several studies in rats have identified QTL for traits potentially related to those studied here. *Tame-1* overlaps with earlier identified QTL influencing several anxiety-related traits (TERENINA-RIGALDIE *et al.* 2003), rearing behavior (FERNANDEZ-TERUEL *et al.* 2002), and adrenal gland weight (SOLBERG *et al.* 2006). *Tame-2* overlaps with two QTL for activity and

anxiety-related behaviors (TERENINA-RIGALDIE *et al.* 2003; CONTI *et al.* 2004). It is reassuring that phenotypes similar to some of those studied here show linkage to similar genomic locations. However, in the absence of information on the molecular basis of these QTL, it cannot presently be determined whether alleles at the same genes are responsible.

**A genetic network for tameness:** Epistasis affects the expression of numerous traits (PHILLIPS 2008). For behavioral quantitative traits, however, epistatic networks identified by genome scans remain rare (for exceptions, see WRIGHT *et al.* 2006; BAILEY *et al.* 2008). In this cross, a network of *Tame-1* and *Tame-2* and three additional loci that were identified only as part of these epistatic pairs influence tameness (Figure 5A).

The additive effect of *Tame-1* was robust across genetic backgrounds (Figure 5), suggesting that *Tame-1* is a major



**TABLE 4**  
**Epistatic pairs of QTL identified at genomewide significance**

Trait	Locus 1			Locus 2			% variance <sup>c</sup>
	Chr <sup>a</sup>	Peak <sup>b</sup>	Position <sup>b</sup>	Chr <sup>a</sup>	Peak <sup>b</sup>	Position <sup>b</sup>	
Tameness (PC1)	1	58	42–96	8	89	44–97	9.6
	1	69	67–75	4	25	21–37	8.7
	1	69	67–71	19	53	47–61	8.7
	6	82	67–94	8	73	67–81	6.3
	8	71	66–80	19	52	43–61	6.8
Flight	1	45	43–48	9	8	0–14	6.9
Boxing	1	75	73–79	8	95	88–97	6
Time spent moving (LDT)	8	86	83–88	9	84	82–84	5.4
Fecal boli (LDT)	1	10	5–10	10	59	49–71	6.6
	4	8	1–15	13	22	17–27	6.1
Startle response	2	48	33–55	10	69	63–86	6.5
Body weight	7	55	22–68	8	97	95–97	6.3
	8	96	94–97	18	16	12–17	6
Adrenal gland weight	3	0	0–2	11	43	38–48	5.2
Corticosterone	5	25	18–35	6	82	73–92	6.2

LDT: light–dark test.

<sup>a</sup>Chromosome.

<sup>b</sup>Centimorgans.

<sup>c</sup>Residual phenotypic variance explained after accounting for fixed effects.

locus influencing tameness. In contrast, *Tame-2* seems to act as a capacitor of other loci. When it is homozygous for the allele from the aggressive line, it magnifies the effects of all three loci it interacts with (Figure 5, C, D, and F). This is reminiscent of epistatic loci underlying growth in chicken (CARLBORG *et al.* 2006). In addition, the effect of *Tame-2* itself is highly dependent on other loci. In a tame background, *i.e.*, where the other loci are homozygous for the allele from the tame line, *Tame-2* has at most a small effect on tameness.

The epistatic network raises interesting questions about the role of *Tame-1* and *Tame-2* during selection for tameness and aggression. Due to the relatively invariant effect of *Tame-1*, it can be selected for in many genetic backgrounds, driving alternative alleles rapidly to fixation. On the other hand, the homozygous aggressive genotype at *Tame-2* might have had an initial role in selection by magnifying the effects of other loci, allowing them to become more prominent targets of selection, while the allele from the tame line would have decreased the response of other loci to selection. The increased frequency of homozygous tame genotypes at loci other than *Tame-2* will, however, decrease the selective advantage of the tame allele at *Tame-2*, due to its small effect in this background. Given the intricate interactions between *Tame-2* and the other loci, it is an intriguing possibility that *Tame-2* might harbor multiple alleles in the current tame and aggressive rat lines. Thus, the single tame and aggressive alleles might in fact be

average effects across several alleles. A more in-depth analysis of patterns of polymorphism at *Tame-2* and other loci might shed light on this.

**White coat color and tameness:** Many domestic animals across a wide range of species are distinguished from their wild relatives by conspicuous coat color variants. Possible explanations include direct selection for coat color variants by humans (*e.g.*, PIELBERG *et al.* 2008) and removal of selective pressures for camouflage. It is further conceivable that coat color variation is a pleiotropic effect of alleles influencing other traits and particularly behavior, including the level of tameness (KEELER and KING 1942; COTTLE and PRICE 1987; HAYSEN 1997).

The F<sub>2</sub> rats provide an excellent opportunity to test whether loci influencing tameness also affect white coat spotting. If the same genes are responsible, F<sub>2</sub> animals carrying coat spots should be more tame than those without. However, this was not observed (see RESULTS). Similarly, the QTL for coat spotting shows no linkage to tameness or any other trait (Figure 4B), and neither *Tame-1* nor *Tame-2* is linked to coat spotting (Figure S10). Hence, we find no evidence for white spotting being caused by the same loci that contribute to tameness. Pleiotropic effects linking tameness and coat color may occur in other species, or even in other lines of rats, but such scenarios are clearly not strengthened by these results.

It is noteworthy that the QTL for white coat spots on chromosome 14 contains at its center the *Kit* gene

(RefSeq NM\_022264), which encodes a tyrosine-kinase receptor involved in melanoblast migration (YOSHIDA *et al.* 2001). Allelic variants of homologs of rat *Kit*, or of the gene for the *Kit ligand* (*Kitl*; RefSeq NM\_021843), are known to cause white coat color variants in mice (JACKSON 1994), pigs (MARKLUND *et al.* 1998), horses (HAASE *et al.* 2007), and stickleback fish (MILLER *et al.* 2007). Thus, *Kit* is an excellent candidate for causing the white coat spots in the rats studied here.

**Conclusions:** We present a genetic analysis of traits associated with tameness in a rat model of animal domestication. Tameness is found to be influenced by two major loci, which are part of a five-locus epistatic network. A possibility not explored here are epistatic interactions involving more than two loci. Such interactions are, however, very difficult to detect given the sample size limitations in mammals.

The confidence intervals for the two tameness loci contain 744 (*Tame-1*) and 339 (*Tame-2*) genes annotated in the Ensembl database, respectively. Since few genes underlying QTL for any behavior have been identified (for two notable exceptions, see YALCIN *et al.* 2004 and WATANABE *et al.* 2007), and none of them are located in *Tame-1* or *Tame-2*, it seems premature to speculate about what genes underlie *Tame-1* and *Tame-2*. Rather, fine-mapping approaches such as advanced intercross lines, as well as other approaches, are needed to clarify what genes are involved.

This work greatly benefited from SNP genotyping courteously provided by Ivo Gut and Marc Lathrop at the Centre National de Genotypage, Genome Institute of the French Atomic Energy Commission, Evry, France. We are indebted to Lysann Wagner and Claudia Nestler for experimental assistance; to Sandra Peter, Petra Madaj-Sterba, and Gudrun Lemm for assistance with animal maintenance; and to Inger Jonasson, Jenny Jonsson, Katarina Davidsson, and Ulf Gyllenstein at the Uppsala Genome Center for help with genotyping and for being great hosts to F.W.A. in Uppsala. We thank two anonymous reviewers for their constructive comments on the manuscript. This work was funded by the Max Planck Society. The work by Ö.C. and F.B. was funded by the Swedish Foundation for Strategic Research and the Swedish Research Council for Environment, Agricultural Sciences and Spatial Planning.

#### LITERATURE CITED

- ALBERT, F. W., O. SHCHEPINA, C. WINTER, H. RÖMPLER, D. TEUPSER *et al.*, 2008 Phenotypic differences in behavior, physiology and neurochemistry between rats selected for tameness and for defensive aggression towards humans. *Horm. Behav.* **53**: 413–421.
- ALVAREZ-CASTRO, J. M., and O. CARLBORG, 2007 A unified model for functional and statistical epistasis and its application in quantitative trait loci analysis. *Genetics* **176**: 1151–1167.
- BAILEY, J. S., L. GRABOWSKI-BOASE, B. M. STEFFY, T. WILTSHIRE, G. A. CHURCHILL *et al.*, 2008 Identification of quantitative trait loci for locomotor activation and anxiety using closely related inbred strains. *Genes Brain Behav.* **7**: 761–769.
- BATES, D., 2007 *lme4: Linear Mixed-Effects Models Using S4 Classes*. R package version 0.99875–9. <http://cran.r-project.org/web/packages/lme4/index.html>.
- BELYAEV, D. K., and P. M. BORODIN, 1982 The influence of stress on variation and its role in evolution. *Biol. Zent. Bl.* **100**: 705–714.
- BESNIER, F., and O. CARLBORG, 2007 A general and efficient method for estimating continuous IBD functions for use in genome scans for QTL. *BMC Bioinformatics* **8**: 440.
- CARLBORG, O., and L. ANDERSSON, 2002 Use of randomization testing to detect multiple epistatic QTLs. *Genet. Res.* **79**: 175–184.
- CARLBORG, Ö., L. ANDERSSON and B. KINGHORN, 2000 The use of a genetic algorithm for simultaneous mapping of multiple interacting quantitative trait loci. *Genetics* **155**: 2003–2010.
- CARLBORG, O., S. KERJE, K. SCHUTZ, L. JACOBSSON, P. JENSEN *et al.*, 2003 A global search reveals epistatic interaction between QTL for early growth in the chicken. *Genome Res.* **13**: 413–421.
- CARLBORG, O., P. M. HOCKING, D. W. BURT and C. S. HALEY, 2004 Simultaneous mapping of epistatic QTL in chickens reveals clusters of QTL pairs with similar genetic effects on growth. *Genet. Res.* **83**: 197–209.
- CARLBORG, O., L. JACOBSSON, P. AHGREN, P. SIEGEL and L. ANDERSSON, 2006 Epistasis and the release of genetic variation during long-term selection. *Nat. Genet.* **38**: 418–420.
- CHURCHILL, G. A., and R. W. DOERGE, 1994 Empirical threshold value for quantitative trait mapping. *Genetics* **138**: 963–971.
- CONTI, L. H., M. JIROUT, L. BREEN, J. J. VANELLA, N. J. SCHORK *et al.*, 2004 Identification of quantitative trait loci for anxiety and locomotion phenotypes in rat recombinant inbred strains. *Behav. Genet.* **34**: 93–103.
- COTTLE, C. A., and E. O. PRICE, 1987 Effects of the nonagouti pelage-color allele on the behavior of captive wild Norway rat (*Rattus norvegicus*). *J. Comp. Psychol.* **101**: 390–394.
- DARWIN, C., 1868 *The Variation of Animals and Plants Under Domestication*, Ed. 1. John Murray, London.
- DIAMOND, J., 2002 Evolution, consequences and future of plant and animal domestication. *Nature* **418**: 700–707.
- DRISCOLL, C. A., M. MENOTTI-RAYMOND, A. L. ROCA, K. HUPE, W. E. JOHNSON *et al.*, 2007 The Near Eastern origin of cat domestication. *Science* **317**: 519–523.
- ERIKSSON, J., G. LARSON, U. GUNNARSSON, B. BED'HOM, M. TIXIER-BOICHARD *et al.*, 2008 Identification of the Yellow skin gene reveals a hybrid origin of the domestic chicken. *PLoS Genet.* **4**: e1000010.
- FERNANDEZ-TERUEL, A., R. M. ESCORIHUELA, J. A. GRAY, R. AGUILAR, L. GIL *et al.*, 2002 A quantitative trait locus influencing anxiety in the laboratory rat. *Genome Res.* **12**: 618–626.
- FLINT, J., 2003 Analysis of quantitative trait loci that influence animal behavior. *J. Neurobiol.* **54**: 46–77.
- GIBBS, R. A., G. M. WEINSTOCK, M. L. METZKER, D. M. MUZYNY, E. J. SODERGREN *et al.*, 2004 Genome sequence of the Brown Norway rat yields insights into mammalian evolution. *Nature* **428**: 493–521.
- GREEN, S., K. FALLS and S. CROOKS, 1990 *Documentation for CRI-MAP v2.4*. Washington University School of Medicine, St. Louis.
- HAASE, B., S. A. BROOKS, A. SCHLUMBAUM, P. J. AZOR, E. BAILEY *et al.*, 2007 Allelic heterogeneity at the equine KIT locus in dominant white (W) horses. *PLoS Genet.* **3**: 2101–2108.
- HALEY, C. S., and S. A. KNOTT, 1992 A simple regression method for mapping quantitative trait loci in line crosses using flanking markers. *Heredity* **69**: 315–324.
- HALEY, C. S., S. A. KNOTT and J.-M. ELSÉN, 1994 Mapping quantitative trait loci in crosses between outbred lines using least-squares. *Genetics* **136**: 1195–1207.
- HARE, B., I. Z. PLYUSNINA, N. IGNACIO, O. SCHEPINA, A. STEPIKA *et al.*, 2005 Social cognitive evolution in captive foxes is a correlated by-product of experimental domestication. *Curr. Biol.* **15**: 226–230.
- HAYSEN, V., 1997 Effects of the nonagouti coat-color allele on behavior of deer mice (*Peromyscus maniculatus*): a comparison with Norway rats (*Rattus norvegicus*). *J. Comp. Psychol.* **111**: 419–423.
- JACKSON, I. J., 1994 Molecular and developmental genetics of mouse coat color. *Annu. Rev. Genet.* **28**: 189–217.
- KALINOWSKI, S. T., M. L. TAPER and T. C. MARSHALL, 2007 Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Mol. Ecol.* **16**: 1099–1106.
- KEELER, C. E., and H. D. KING, 1942 Multiple effects of coat color genes in the Norway rat, with special reference to temperament and domestication. *J. Comp. Psychol.* **34**: 241–250.
- KRUK, M. R., J. HALÁSZ, W. MEELIS and J. HALLER, 2004 Fast positive feedback between the adrenocortical stress response and a brain mechanism involved in aggressive behavior. *Behav. Neurosci.* **118**: 1062–1070.

- LARSON, G., K. DOBNEY, U. ALBARELLA, M. FANG, E. MATISOO-SMITH *et al.*, 2005 Worldwide phylogeography of the wild boar reveals multiple centers of pig domestication. *Science* **307**: 1618–1621.
- LE ROUZIC, A., and J. M. ALVAREZ-CASTRO, 2008 Estimation of genetic effects and genotype-phenotype maps. *Evol. Bioinform. Online* **4**: 225–235.
- LYNCH, M., and B. WALSH, 1998 *Genetics and Analysis of Quantitative Traits*. Sinauer Associates, Sunderland, MA.
- MARKLUND, S., J. KIJAS, H. RODRIGUEZ-MARTINEZ, L. RONNSTRAND, K. FUNA *et al.*, 1998 Molecular basis for the dominant white phenotype in the domestic pig. *Genome Res.* **8**: 826–833.
- MILLER, C. T., S. BELEZA, A. A. POLLEN, D. SCHLUTER, R. A. KITTLES *et al.*, 2007 cis-regulatory changes in kit ligand expression and parallel evolution of pigmentation in sticklebacks and humans. *Cell* **131**: 1179–1189.
- NADERI, S., H.-R. REZAEI, F. POMPANON, M. G. B. BLUM, R. NEGRINI *et al.*, 2008 The goat domestication process inferred from large-scale mitochondrial DNA analysis of wild and domestic individuals. *Proc. Natl. Acad. Sci. USA* **105**: 17659–17664.
- NAUMENKO, E. V., N. K. POPOVA, E. M. NIKULINA, N. N. DYGALO, G. T. SHISHKINA *et al.*, 1989 Behavior, adrenocortical activity, and brain monoamines in norway rats selected for reduced aggressiveness towards man. *Pharmacol. Biochem. Behav.* **33**: 85–91.
- O'CONNELL, J. R., and D. E. WEEKS, 1998 PedCheck: a program for identification of genotype incompatibilities in linkage analysis. *Am. J. Hum. Genet.* **63**: 259–266.
- PEREZ-ENCISO, M., and I. MISZTAL, 2004 Qxpak: a versatile mixed model application for genetical genomics and QTL analyses. *Bioinformatics* **20**: 2792–2798.
- PHILLIPS, P. C., 2008 Epistasis—the essential role of gene interactions in the structure and evolution of genetic systems. *Nat. Rev. Genet.* **9**: 855–867.
- PIELBERG, G. R., A. GOLOVKO, E. SUNDSTROM, I. CURIK, J. LENNARTSSON *et al.*, 2008 A cis-acting regulatory mutation causes premature hair graying and susceptibility to melanoma in the horse. *Nat. Genet.* **40**: 1004–1009.
- PINHEIRO, J., D. BATES, S. DEBROY, D. SARKAR and THE R CORE TEAM, 2008 *nlme: Linear and Nonlinear Mixed Effects Models*. R package version 3.1–87. <http://cran.r-project.org/web/packages/nlme/index.html>.
- PLYUSNINA, I. Z., and I. OSKINA, 1997 Behavioral and adrenocortical responses to open-field test in rats selected for reduced aggressiveness toward humans. *Physiol. Behav.* **61**: 381–385.
- PONG-WONG, R., A. W. GEORGE, J. A. WOOLLIAMS and C. S. HALEY, 2001 A simple and rapid method for calculating identity-by-descent matrices using multiple markers. *Genet. Sel. Evol.* **33**: 453–471.
- POPOVA, N. K., V. S. NAUMENKO, I. Z. PLYUSNINA and A. V. KULIKOV, 2005 Reduction in 5-HT1A receptor density, 5-HT1A mRNA expression and functional correlates for 5-HT1A receptors in genetically defined aggressive rats. *J. Neurosci. Res.* **80**: 286–292.
- R DEVELOPMENT CORE TEAM, 2008 *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna.
- SAAR, K., A. BECK, M. T. BIHOREAU, E. BIRNEY, D. BROCKLEBANK *et al.*, 2008 SNP and haplotype mapping for genetic analysis in the rat. *Nat. Genet.* **40**: 560–566.
- SAVOLAINEN, P., Y.-P. ZHANG, J. LUO, J. LUNDEBERG and T. LEITNER, 2002 Genetic evidence for an East Asian origin of domestic dogs. *Science* **298**: 1610–1613.
- SCHUELKE, M., 2000 An economic method for the fluorescent labeling of PCR fragments. *Nat. Biotechnol.* **18**: 233–234.
- SOLBERG, L. C., A. E. BAUM, N. AHMADIYEH, K. SHIMOMURA, R. H. LI *et al.*, 2006 Genetic analysis of the stress-responsive adrenocortical axis. *Physiol. Genomics* **27**: 362–369.
- SUMMERS, C. H., M. J. WATT, T. L. LING, G. L. FORSTER, R. E. CARPENTER *et al.*, 2005 Glucocorticoid interaction with aggression in non-mammalian vertebrates: reciprocal action. *Eur. J. Pharmacol.* **526**: 21–35.
- TERENINA-RIGALDIE, E., M. P. MOISAN, A. COLAS, F. BEAUGE, K. V. SHAH *et al.*, 2003 Genetics of behaviour: phenotypic and molecular study of rats derived from high- and low-alcohol consuming lines. *Pharmacogenetics* **13**: 543–554.
- TROY, C. S., D. E. MACHUGH, J. F. BAILEY, D. A. MAGEE, R. T. LOFTUS *et al.*, 2001 Genetic evidence for Near-Eastern origins of European cattle. *Nature* **410**: 1088–1091.
- TRUT, L. N., 1999 Early canid domestication: the farm-fox experiment. *Am. Sci.* **87**: 160–169.
- TRUT, L. N., I. Z. PLYUSNINA and L. A. PRASOLOVA, 2000 Mutations hooded and nonagouti in the Norway rat *Rattus norvegicus*: effects of selection for behavior and photoperiod. *Russ. J. Genet.* **36**: 668–676.
- VILA, C., J. A. LEONARD, A. GOTHERSTROM, S. MARKLUND, K. SANDBERG *et al.*, 2001 Widespread origins of domestic horse lineages. *Science* **291**: 474–477.
- WATANABE, A., T. TOYOTA, Y. OWADA, T. HAYASHI, Y. IWAYAMA *et al.*, 2007 Fabp7 maps to a quantitative trait locus for a schizophrenia endophenotype. *PLoS Biol.* **5**: 2469–2483.
- WRIGHT, D., R. K. BUTLIN and Ö. CARLBORG, 2006 Epistatic regulation of behavioural and morphological traits in the zebrafish (*Danio rerio*). *Behav. Genet.* **36**: 914–922.
- YALCIN, B., S. A. G. WILLIS-OWEN, J. FULLERTON, A. MEESAQ, R. M. DEACON *et al.*, 2004 Genetic dissection of a behavioral quantitative trait locus shows that *Rgs2* modulates anxiety in mice. *Nat. Genet.* **36**: 1197–1202.
- YOSHIDA, H., T. KUNISADA, T. GRIMM, E. K. NISHIMURA, E. NISHIOKA *et al.*, 2001 Review: melanocyte migration and survival controlled by SCF/c-kit expression. *J. Invest. Dermatol. Symp. Proc.* **6**: 1–5.

Communicating editor: A. D. LONG



# GENETICS

## Supporting Information

<http://www.genetics.org/cgi/content/full/genetics.109.102186/DC1>

### **Genetic Architecture of Tameness in a Rat Model of Animal Domestication**

**Frank W. Albert, Örjan Carlborg, Irina Plyusnina, Francois Besnier, Daniela Hedwig,  
Susann Lautenschläger, Doreen Lorenz, Jenny McIntosh, Christof Neumann,  
Henning Richter, Claudia Zeising, Rimma Kozhemyakina, Olesya Shchepina,  
Jürgen Kratzsch, Lyudmila Trut, Daniel Teupser, Joachim Thiery, Torsten Schöneberg,  
Leif Andersson and Svante Pääbo**

Copyright © 2009 by the Genetics Society of America

DOI: 10.1534/genetics.109.102186

**FILE S1****Materials and Methods**

**Mapping pedigree:** The pedigree described in this study was initiated from four unrelated tame and four aggressive individuals (two females each; the “F<sub>1</sub>” or “F minus one” generation) mated within line to yield 11 (five tame and six aggressive rats, one male each) F<sub>0</sub> animals. The F<sub>1</sub> animals did not have common parents and at most one common grandparent. The F<sub>0</sub> animals were crossed reciprocally, i.e. both sexes were used from each line. Two out of four tame females and two out of five aggressive females were full siblings of the respective F<sub>0</sub> founder males. Six hybrid F<sub>1</sub> males (three sired by the tame, and three sired by the aggressive F<sub>0</sub> male) were repeatedly mated to 37 F<sub>1</sub> females to produce a total of 733 F<sub>2</sub> rats (383 females). F<sub>1</sub> animals sired by the aggressive F<sub>0</sub> male were mated only to F<sub>1</sub> animals sired by the tame F<sub>0</sub> male, and vice versa.

**Behavioral testing:** A list of all traits we collected can be found in Table 1. At six to eight weeks of age, F<sub>2</sub> rats performed the “glove test”, which measures an animal’s level of tameness/aggression by confronting it with an approaching human hand and attempts at handling (see (ALBERT *et al.* 2008) for details on the testing procedure). Within two weeks, but at least after a four-day break, rats performed an open-field test, followed three days later by a light-dark test. These tests provide various measures of exploratory and anxiety-related behavior. After nine days, rats performed a startle response test, which measures the behavioral response to a sudden acoustic stimulus. 470 (64%) of the F<sub>2</sub> rats then performed a second glove test trial after maximally two weeks. F<sub>1</sub> rats were tested once in the glove test at 12 – 14 weeks of age, and then followed the testing schedule described in (ALBERT *et al.* 2008). All tests were performed with minimal handling following the procedures described in (ALBERT *et al.* 2008). Open field, light-dark and startle response test data were recorded using automated measuring technology (TSE Systems, Bad Homburg, Germany). Glove test trials were videotaped and later analyzed by two independent observers (5% of trials only by one observer) using the software “Interact” (Mangold Software, Arstorf, Germany). Experimenters and observers were blind to the animals’ identity and to further data processing. A set of 11 behaviors (e.g. “attack” or “tolerate handling”) were scored (see (ALBERT *et al.* 2008) for detailed descriptions of the behaviors), and each converted to a numeric measure (e.g. “number of occurrences” or “duration in seconds”; Table 1).

**Serological phenotypes:** We measured a series of serum traits, most of which had earlier been found (ALBERT *et al.* 2008) to differ between the tame and the aggressive rats (Table 1). Serum was analyzed in 684 F<sub>2</sub> rats (357 females) as described in the main text. Corticosterone levels reflect the rats’ response to handling and sacrifice, rather than baseline (undisturbed) levels. This is consistent with their high values compared to basal measures reported in the literature (VAHL *et al.* 2005). Twenty-two F<sub>2</sub> individuals were excluded from further analysis of all serum traits because they appeared to be outliers with respect to the remaining F<sub>2</sub> population: two individuals with chloride values < 90 mmol / l and 16 individuals with chloride values > 110 mmol / l, and four individuals with triglyceride levels > 4 mmol / l.

**Glove test analyses:** To summarize a rat's behavior in the glove test, we performed principal component analysis (PCA) on the individual glove test measures. The resulting principal component (PC) scores represent a linear, weighted combination of individual behaviors. Combining animals from several generations when performing PCA could potentially distort the results. Hence, we used only F<sub>2</sub> animals in the analysis.

F<sub>0</sub> and F<sub>1</sub> animals performed only one glove test trial and were scored by a single observer. For comparing F<sub>2</sub> animals' glove tests to those of the F<sub>0</sub> and F<sub>1</sub> generations, we used only observations from the first trial, and from the same observer who scored the F<sub>0</sub> and F<sub>1</sub> animals. We applied the PCA regression coefficients obtained in F<sub>2</sub> animals to the single observations from F<sub>0</sub>, F<sub>1</sub>, and the comparable subset of single F<sub>2</sub> observations. A small number of F<sub>2</sub> animals had PC scores that exceeded the range defined by the most tame and the most aggressive F<sub>0</sub> animals. Since QTL mapping can be sensitive to phenotypic outliers, we set these extreme individuals' phenotypes to the most extreme values observed in F<sub>0</sub> animals. Adjusting outliers this way did not greatly alter the mapping results.

**Marker ascertainment and genotyping procedures:** Animals were genotyped with 152 microsatellite and 49 single nucleotide polymorphism (SNP) markers. SNPs were added after a preliminary scan for QTLs using only the microsatellite data (not shown), to improve genome coverage and to obtain more data in putative QTL regions. Microsatellite markers were ascertained from public databases. SNP markers were ascertained from a panel of 16,927 SNPs under study at the CEA/IG-Centre National de Genotypage, Evry, France. All markers were selected based on their segregation patterns in the outbred tame and aggressive rat lines as determined from preliminary genotyping of a panel of F<sub>0</sub> animals. All markers used in the QTL mapping are listed in Table S2. DNA was isolated either from lung tissue according to the NucPrep Protocol for animal tissues (Applied Biosystems, Foster City, CA, USA) or from spleen tissue using the DNeasy Blood & Tissue Kit (Quiagen GmbH, Hilden, Germany) according to the manual's section "Purification of Total DNA from Animal Tissues (Spin-Column Protocol)". The extraction protocols were slightly modified. For NucPrep, digestion at 65 °C for 1 h was preceded by overnight incubation of the tissue samples at room temperature in NucPrep Digestion Buffer and Proteinase K (both Applied Biosystems), and the pre-filtration step was skipped. For DNeasy, two final elution steps were performed, each with 100 µl of Buffer AE. For all markers, polymerase chain reaction (PCR) was performed in 384-well format. Microsatellite marker PCRs were performed using the M13-primer PCR system (SCHUELKE 2000). 15 ng of dried DNA per sample were used in a 10 µl PCR mix containing (per sample) 1.0 µl of PCR buffer (Quiagen), 0.5 µl DMSO (Merck KG, Darmstadt, Germany), 0.4 µl dNTPs (5 mM) (Amersham Biosciences, Buckinghamshire, England), 0.2 µl forward and reverse primer in a 1 / 10 ratio (1 / 10 µM) (Thermo Electron, Ulm, Germany), 0.2 µl M13-tail (10 µM) labeled with one of three fluorophores [6-fam (Thermo Electron), Ned or Vic (both Applied Biosystems)], 7.66 µl of water and 0.039 µl of HotStarTaq® Plus DNA polymerase (5 U / µl) (Quiagen). The M13-tail used in all reactions was 5'-[fluorophore]-CACGACGTTGTAAAACGAC. PCR was performed on Eppendorf Mastercycler ep384 thermocyclers (Eppendorf, Hamburg, Germany) according to the following touch-down PCR protocol: 95 °C for 5 min., 44 cycles of each at 94 °C for 30 s, a variable temperature for 30 s



and 72 °C for 30 s. The variable temperature was 65 °C in the first cycle and decreased by 1 °C in the following 5 cycles, then decreased by 0.5 °C for 17 cycles and then held constant at 52 °C for the remaining 21 cycles. PCR products were held at 8 °C after program completion. Up to 10 PCR products of different expected lengths and labeled with different dyes were pooled and analyzed on an ABI3730 DNA Analyzer (Applied Biosystems). Microsatellite genotypes were determined using the software GeneMapper Version 4.0 (Applied Biosystems) and all genotypes were manually double-checked. SNP assays were ordered from ABI Applied Biosystems. PCR cocktails containing 10 ng of dried DNA / sample were prepared according to the assay manufacturer's specifications, but using "ABgene Absolute QPCR Rox Mix X2" (Applied Biosystems) instead of "TaqMan Universal PCR Master Mix". PCR was performed on Eppendorf Mastercycler ep384 thermocyclers (Eppendorf) as follows: 10 min. at 95 °C, followed by 40 cycles of 15 s at 92 °C and 1 min. at 60 °C; reactions were held at 8 °C after program completion. Intensity scans of PCR products were performed on an ABI 7900HT Sequence Detection System v2.2 (Applied Biosystems). SNP genotypes were called automatically as part of the scanning process, and genotype plots inspected visually. Preliminary genotyping of 16,927 SNPs had been performed using the Illumina GoldenGate assay according to manufacturer's instruction in an Illumina Beadlab, as described (SAAR *et al.* 2008).

**Details on linkage map construction:** We constructed sex-averaged linkage maps for each chromosome using a version of the program *crimap* (GREEN *et al.* 1990) modified to handle large pedigrees by the University of California Davis, Veterinary Genetics Laboratory. We used the option *FIXED* to estimate marker distances in centiMorgan (cM) on markers ordered according to their physical chromosomal locations as annotated in the UCSC Genome Browser. Marker order was confirmed using *crimap*'s *FLIPS* option. Markers that were not annotated in the genome browser were placed on the linkage map using the option *ALL*. We used the option *CHROMPIC* to identify inferred double recombination events. For all such events, genotype data was double-checked for errors.

**Computation of linkage map information content:** We estimated the information content at marker positions from the respective gametic identical-by-decent (IBD) matrices  $X_g$  (see below for the matrix estimation procedure). For a given position, the IBD matrices specify which individuals have inherited identical alleles. At informative positions, many allele pairs can be assigned unambiguously (IBD = 1 or IBD = 0). At less informative positions, some estimated IBD estimates will

tend towards 0.5. We estimated the information content at a position as  $2 \frac{1}{n} \sum |0.5 - X_g|$ , where  $n$  is the number of alleles in the pedigree. The information content statistic takes a value of one for perfectly informative markers, and zero if no allele can be traced to its pedigree origin. On the X-chromosome, we calculated a lower limit for the information content at marker positions as the average difference in allele frequency between tame and aggressive  $F_{-1}$  founders, weighted by the respective allele frequency across all  $F_{-1}$  founders.

**Haplotype-based inference of missing genotypes:** Missing  $F_0$  genotypes can lead to a loss of power in QTL mapping because some alleles in  $F_2$  animals might not be traced reliably to parental lines. We sought to overcome this limitation by including the genotyped parents of  $F_0$  animals (the  $F_{-1}$ ) in the analysis, and by tracing alleles back to them. This way, we sought to infer missing genotypes of  $F_0$  and  $F_1$  animals based on their ancestors' and offspring genotypes. We first determined the phase of informative markers for all individuals in the pedigree. Then, we used a genetic algorithm to iteratively generate chromosome segments carrying putative haplotypes. For each such segment, the algorithm calculates the likelihood that the segment is compatible with the phase of the informative markers, and keeps the most likely haplotype from each iteration until it converges to an optimum. We then inferred the missing genotype information in  $F_0$  and  $F_1$  animals from the haplotypes of related  $F_{-1}$  and  $F_2$  individuals. Details of this procedure for inference of haplo- and genotypes will be published elsewhere.

**Single QTL mapping on sex chromosomes:** The analyses described thus far were designed to handle autosomal data. To analyze the X-chromosome, we used the software QxPak (PEREZ-ENCISO and MISZTAL 2004), which is able to handle sex chromosomal data appropriately. For each trait, we first fitted an additive QTL using the option 'fix\_a'. Where this yielded a significant ( $p < 0.005$ ) QTL, we then tested for dominance using option 'fix\_ad'. Since the permutation-based significance thresholds derived for the autosomes cannot be directly applied to the X chromosome, we assumed QTLs with a nominal p-value  $< 0.001$  ( $0.005$ ) to be significant (suggestive) at a genome-wide level, as suggested in the QxPak manual.

To test for linkage to the Y chromosome, we note that  $F_2$  males in our pedigree carry one of only two Y chromosomes, one derived from the tame  $F_0$  male, one from the aggressive  $F_0$  male. Due to the design of our cross,  $F_2$  males carrying the Y chromosome from one parental line always carry mitochondria (and potentially other maternally inherited factors) from the same line. However, while  $F_2$  females share such maternal factors with  $F_2$  males, they do not carry the Y chromosomes. This allows us to disentangle the effect of the Y chromosome by comparing the phenotypes of  $F_2$  males carrying the Y chromosome inherited from the tame  $F_0$  male, of males carrying the Y chromosome inherited from the aggressive  $F_0$  male, and of  $F_2$  females that were sired by the respective  $F_1$  males. A Y-linked locus should lead to a phenotype difference between the two groups of  $F_2$  males, while the two groups of  $F_2$  females should not differ. In an analysis of variance (ANOVA), we tested for interactions between the sex of  $F_2$  animals and the strain origin of the Y chromosome carried by their fathers. Covariates were included in the ANOVA as listed in Table S6. We assumed linkage to the Y chromosome if the interaction term was significant ( $p < 0.005$ ) and if a post-hoc T-test showed a significant ( $p < 0.05$ ) difference between  $F_2$  males, but not between  $F_2$  females.

- ALBERT, F. W., O. SHCHEPINA, C. WINTER, H. RÖMPLER, D. TEUPSER *et al.*, 2008 Phenotypic differences in behavior, physiology and neurochemistry between rats selected for tameness and for defensive aggression towards humans. *Hormones and Behavior* **53**: 413-421.
- BESNIER, F., and O. CARLBORG, 2007 A general and efficient method for estimating continuous IBD functions for use in genome scans for QTL. *Bmc Bioinformatics* **8**.
- CARLBORG, O., and L. ANDERSSON, 2002 Use of randomization testing to detect multiple epistatic QTLs. *Genetical Research* **79**: 175-184.
- CARLBORG, Ö., L. ANDERSSON and B. KINGHORN, 2000 The Use of a Genetic Algorithm for Simultaneous Mapping of Multiple Interacting Quantitative Trait Loci. *Genetics* **155**: 2003-2010.
- CHURCHILL, G. A., and R. W. DOERGE, 1994 Empirical threshold value for quantitative trait mapping. *Genetics* **138**: 963-971.
- GREEN, S., K. FALLS and S. CROOKS, 1990 *Documentation for CRI-MAP v2.4*. Washington University School of Medicine, St. Louis.
- HALEY, C. S., and S. A. KNOTT, 1992 A Simple Regression Method For Mapping Quantitative Trait Loci In Line Crosses Using Flanking Markers. *Heredity* **69**: 315-324.
- HALEY, C. S., S. A. KNOTT and J.-M. ELSEN, 1994 Mapping Quantitative Trait Loci In Crosses Between Outbred Lines Using Least-Squares. *Genetics* **136**: 1195-1207.
- PEREZ-ENCISO, M., and I. MISZTAL, 2004 Qxpak: a versatile mixed model application for genetical genomics and QTL analyses. *Bioinformatics* **20**: 2792-2798.
- PONG-WONG, R., A. W. GEORGE, J. A. WOOLLIAMS and C. S. HALEY, 2001 A simple and rapid method for calculating identity-by-descent matrices using multiple markers. *Genetics Selection Evolution* **33**: 453-471.
- SAAR, K., A. BECK, M. T. BIHOREAU, E. BIRNEY, D. BROCKLEBANK *et al.*, 2008 SNP and haplotype mapping for genetic analysis in the rat. *Nature Genetics* **40**: 560-566.
- SCHUELKE, M., 2000 An economic method for the fluorescent labeling of PCR fragments. *Nature Biotechnology* **18**: 233-234.
- VAHL, T. P., Y. M. ULRICH-LAI, M. M. OSTRANDER, C. M. DOLGAS, E. E. ELFERS *et al.*, 2005 Comparative analysis of ACTH and corticosterone sampling methods in rats. *American Journal of Physiology: Endocrinology and Metabolism* **289**: E823-E828.

**FILE S2**

A compressed folder is available at <http://www.genetics.org/cgi/content/full/genetics.109.102186/DC1>.

This folder contains four text files that contain raw data (as described below). This description is also available in the **Read Me.txt** file located in the compressed folder.

1. "genotypes.txt" – contains genotypes and marker info
  - first line contains marker names
  - second line contains markers' chromosomes
  - third line contains markers' genetic position on a given chromosome
  - the rest of the file contains the genotypes:
    - first column is animal id
    - there are *\*two\** columns per marker, one for each allele
    - allele codes are arbitrary numbers, and do not by themselves contain information about strain origin or the genotype
    - the two allele columns per marker do *\*not\** indicate whether an allele was inherited from the mother or the father
    - missing genotypes are coded as '0'
    - on the X-chromosome, '9' indicates the "missing" allele in hemizygous males
2. "pedigree\_fixed\_effects\_covariates.txt"
  - first column is animal id
  - second column is an animal's father ('0' for the first generation)
  - third column: is an animal's mother ('0' for the first generation)
  - fourth column is sex: males are '1', females are '2'
  - fifth column indicates an animals strain (tame or aggressive) or the respective generation in the cross (F1 or F2)
  - remaining columns are numerical covariates (measured only in F2 animals)
3. "phenotypes.txt"
  - first column is animal id
  - only F2 animals were measured (beginning with animal ID 63), and only F2s are listed in this file
  - with the exception of glove test traits, all traits are "raw", i.e. not adjusted for family effects or covariates
  - glove test behaviors in this file *\*are\** adjusted for observer, trial and family effects - see "raw\_glove\_data.txt" for raw glove test data
4. "raw\_glove\_data.txt"
  - data from the glove test, directly as scored from the video tapes
  - this data forms the basis for the principal components used to calculate tameness and aggression scores
  - only F2 animals are listed (beginning with animal ID 63)






FIGURE S1.—Flow diagram of the search strategy for single QTLs and epistatic QTL pairs.




FIGURE S2.—Phenotype ranges for traits recorded in the glove test. Solid boxes represent the 50% of trait values closest to the median (bold line). Whiskers represent the low and high 25% of trait values.




FIGURE S3.—Phenotype ranges for traits recorded in the open field, light-dark and startle response tests. Solid boxes represent the 50% of trait values closest to the median (bold line). Whiskers represent the low and high 25% of trait values. For clarity, potential outliers are not shown.




FIGURE S4.—Phenotype ranges for body and organ weights. Solid boxes represent the 50% of trait values closest to the median (bold line). Whiskers represent the low and high 25% of trait values. For clarity, potential outliers are not shown.




FIGURE S5.—Phenotype ranges for serum traits. Solid boxes represent the 50% of trait values closest to the median (bold line). Whiskers represent the low and high 25% of trait values. For clarity, potential outliers are not shown.






FIGURE S6.—Phenotype distributions for traits recorded in the glove test. Red: female F2 rats, blue: male F2 rats




FIGURE S7.—Phenotype distributions for traits recorded in the open field, light-dark, and startle response tests. Red: female F2 rats, blue: male F2 rats.




FIGURE S8.—Phenotype distributions for body and organ weights. Organ weights are shown relative to body weight. Red: female F2 rats, blue: male F2 rats




FIGURE S9.—Phenotype distributions for serum traits. Red: female F2 rats, blue: male F2 rats




FIGURE S10.—QTLs for tameness do not influence white coat spotting. Red line: tameness. Black line: spotting. Black upward tickmarks: microsatellite marker positions. Red tickmarks: SNP marker positions



**TABLE S1**  
**Phenotype modelling**

Phenotype group	Fixed effects	Random effects	Covariates	Adjusted effects	Effects included in QTL model
Glove test*	sex, trial	litter, observer	-	litter, trial, observer	sex
Open field & light-dark tests	sex	litter	-	litter	sex
Startle response	sex	litter	body weight	litter	sex, body weight
Organ weights	sex	litter	body weight	litter	sex, body weight
Body weight	sex	litter	age	litter	sex, age
Serum phenotypes	sex	litter	age <sup>§</sup>	litter, batch	sex, age <sup>§</sup>

To account for repeated observations of the same rat, trial and observer were modeled as being nested within individual rats. <sup>§</sup> age was included for traits for which a significant correlation (Pearson's product momentum correlation,  $p < 0.05$ ) between trait and age was found (ALAT, ASAT, AP, creatinine, urea, chloride, iron, protein, albumin and fT4, but not for glucose, calcium, fT3 and corticosterone).

**TABLE S2****Genetic markers used in QTL mapping**

Table S2 is available as an Excel file at <http://www.genetics.org/cgi/content/full/genetics.109.102186/DC1>.

**TABLE S3**  
**Sex differences**

Trait	p-value	females		males		Unit
		mean	s.e.m. <sup>1</sup>	mean	s.e.m. <sup>1</sup>	
<b>Open Field Test</b>						
time spent in center	0.4	3.46	0.2	3.43	0.2	%
time spent in corner	0.06	0.65	0.01	0.62	0.01	%
time spent moving	< 0.0001	41	0.5	36	0.6	%
time spent rearing	< 0.0001	23	0.7	17	0.6	s
locomotion speed	< 0.0001	29	0.4	25	0.4	cm / s
fecal boli	0.6	2.1	0.1	2.2	0.1	count
<b>Light - Dark Test</b>						
time spent in light compartment	0.6	26	0.9	26	1.1	%
time spent moving	< 0.0001	52	0.5	48	0.6	%
time spent rearing	< 0.0001	22	0.6	18	0.6	s
locomotion speed in light compartment	< 0.0001	22	0.5	19	0.6	cm / s
fecal boli	0.01	1.9	0.1	2.4	0.1	count
<b>Startle Response Test</b>						
startle response	0.01	76	9.9	72	16.4	g
<b>Anatomy</b>						
body weight	< 0.0001	111	1.3	222	2.2	g
adrenal gland weight	< 0.0001	42	0.4	40	0.4	mg
heart weight	< 0.0001	0.33	0.008	0.38	0.009	g
kidney weight	< 0.0001	0.04	0.004	0.07	0.007	g
lung weight	< 0.0001	0.67	0.01	0.64	0.01	g
spleen weight	< 0.0001	0.16	0.004	0.15	0.005	g
<b>Serum traits</b>						
<i>Hormones</i>						
corticosterone	< 0.0001	1650	26	1192	25	ng / ml
fT3	< 0.0001	5.3	0.05	4.8	0.05	pmol / l
fT4	< 0.0001	38.5	0.4	45.3	0.4	pmol / l
<i>Enzymes</i>						
ALAT	< 0.0001	0.49	0.006	0.6	0.008	μkat / l
AP	< 0.0001	1.5	0.01	2	0.02	μkat / l
ASAT	< 0.0001	2.7	0.05	2.3	0.03	μkat / l
<i>Substrates</i>						
creatinine	< 0.0001	36	0.3	33	0.3	μkat / l
glucose	< 0.0001	7.00	0.1	10.5	0.2	μmol / l

urea	< 0.0001	6.2	0.06	5.3	0.05	mmol / l
<i>Serum protein</i>						
total protein	0.014	71	0.2	70	0.2	g / l
albumin	0.0003	48.3	0.2	47.5	0.1	g / l
<i>Electrolytes</i>						
Ca <sup>2+</sup>	< 0.0001	2.82	0.008	2.89	0.007	mmol / l
Cl <sup>-</sup>	< 0.0001	102.5	0.3	101.3	0.2	mmol / l
Fe <sup>2+/3+</sup>	< 0.0001	57	0.7	30	0.4	μmol / l

---

<sup>1</sup> standard error of the mean

**TABLE S4****Correlations between phenotypes**

Table S4 is available as an Excel file at <http://www.genetics.org/cgi/content/full/genetics.109.102186/DC1>.



**TABLE S5**  
**Suggestive autosomal QTLs**

Trait	unit for genetic effects	Chromosome	peak position (cM)	1 LOD support interval (cM)	F-value	additive effect	dominance effect	% residual phenotypic variance explained	females?	males?
<i>Glove Test</i>										
Tameness										
(PC1)	-	3	109	93 - 117	5.4	0.12	0.73	1.4	-	-
Tameness										
(PC1)	-	10	42	11 - 69	5.4	0.4	-0.08	1.4	suggestive	-
Tameness										
(PC1)	-	12	5	2 - 9	5.4	0.13	-0.69	1.4	-	-
Tameness										
(PC1)	-	19	61	45 - 62	5	0.14	-0.75	1.3	-	-
PC2	-	14	5	0 - 25	6	0.32	-0.12	1.7	-	suggestive
PC2	-	11	49	36 - 54	4.8	-0.1	0.46	1.4	-	-
PC3	-	19	15	7 - 25	6	0.2	-0.26	1.7	-	-
PC3	-	10	59	41 - 67	5.1	-0.16	0.39	1.5	-	-
PC3	-	17	0	0 - 16	4.9	0.2	0.26	1.4	suggestive	-
approach	count	3	52	46 - 63	6.8	0.04	-0.15	1.9	suggestive	-
attack	count	8	72	67 - 90	6.6	-0.42	-0.19	1.8	-	-
attack	count	12	6	3 - 10	5.9	-0.18	0.68	1.6	-	-
attack	count	17	15	4 - 24	5.8	0.05	0.83	1.6	-	-
escape	count	10	94	89 - 94	7.8	-0.03	0.52	2.2	-	-
escape	count	18	24	16 - 38	6	-0.09	0.33	1.7	-	-
flight	count	10	49	36 - 71	6.8	-0.33	-0.16	1.9	suggestive	-
flight	count	9	27	16 - 41	5.7	-0.26	0.75	1.6	-	-
move & leave	count	7	1	0 - 5	6.3	0.01	0.38	1.8	-	-
move & leave	count	15	25	21 - 46	6	0.09	-0.36	1.7	-	-
move & leave	count	19	1	0 - 15	5.3	-0.19	-0.15	1.5	-	-
move & leave	count	11	43	33 - 52	5	0.03	-0.36	1.4	-	-
squeak	count	8	97	96 - 97	7.5	-0.02	-5	2.1	-	suggestive
squeak	count	17	1	0 - 11	6.8	1.1	1.5	1.9	-	-
squeak	count	3	63	51 - 72	5.5	-1.2	-0.5	1.6	-	-
boxing	s	6	15	6 - 26	7.3	-0.5	-0.3	2	-	-
boxing	s	1	78	72 - 102	7.1	-0.4	-0.5	2	-	-
boxing	s	19	54	42 - 62	6.8	-0.2	1.1	1.9	suggestive	-

screaming	s	8	72	63 - 82	8.6	-0.73	-0.46	2.3	-	-
screaming	s	19	60	42 - 62	5.4	0.14	1.1	1.4	suggestive	-
screaming	s	10	15	6 - 60	5.3	-0.65	0.17	1.4	-	-
screaming	s	12	8	2 - 23	5.1	-0.4	0.76	1.4	-	-
freezing	s	10	84	68 - 91	6.2	2	-0.7	1.7	-	suggestive
tolerate										
touch	s	18	30	23 - 39	7.2	0.35	-0.22	1.9	significant	-
tolerate										
touch	s	6	86	73 - 93	6.6	1.05	-1.09	1.8	-	-
tolerate										
touch	s	8	0	0 - 17	5.5	0.46	-0.21	1.5	-	-
tolerate										
touch	s	8	80	67 - 92	5.2	0.78	0.99	1.5	-	significant
tolerate										
handling	s	19	17	0 - 27	7	0.79	-0.17	2	-	-
tolerate										
handling	s	14	11	0 - 27	5.6	0.83	0.25	1.6	-	-
tolerate										
handling	s	6	61	51 - 80	5.3	0.68	-0.17	1.5	-	suggestive
tolerate										
handling	s	13	52	32 - 52	5.2	0.46	1.2	1.5	-	-
tolerate										
handling	s	11	49	40 - 54	5	-0.22	1.3	1.4	-	-
<i>Open field</i>										
time spent										
in center	%	8	27	18 - 56	7.8	0.0095	-0.22	2.2	-	suggestive
time spent										
in center	%	10	71	43 - 81	5.9	0.0079	0.0046	1.7	-	-
time spent										
moving	%	2	103	96 - 118	6.3	0.011	0.028	1.8	-	-
time spent										
moving	%	8	94	85 - 97	6.3	0.024	0.004	1.8	-	-
time spent										
moving	%	1	70	62 - 87	6.1	0.021	0.012	1.8	-	-
rearing	s	8	93	85 - 97	4.9	2.3	-0.2	1.4	-	-
locomotion										
speed	cm / s	2	86	83 - 95	6.9	0.5	2.7	2	-	suggestive
locomotion										
speed	cm / s	13	52	48 - 52	5.1	-1.1	-2.4	1.5	-	-
fecal boli	count	2	63	36 - 79	7.6	0.5	-0.2	2.2	suggestive	-
fecal boli	count	9	0	0 - 10	6.6	-0.5	-0.04	1.9	-	suggestive
fecal boli	count	10	80	68 - 91	6.6	0.6	-0.3	1.9	-	-
fecal boli	count	17	18	6 - 25	6.3	0.4	0.6	1.8	-	-

*Light Dark Test*

time spent											
in light	%	6	38	27 - 54	7.6	0.044	-0.013	2.2	-		suggestive
time spent											
in light	%	19	53	43 - 62	7	-0.028	-0.097	2	significant		-
time spent											
in light	%	18	0	0 - 23	6	0.035	-0.025	1.7	-		suggestive
time spent											
in light	%	2	82	72 - 86	5.9	0.0009	0.064	1.7	-		-
time spent											
moving	%	3	0	0 - 2	7.1	0.015	0.036	2	-		-
time spent											
moving	%	7	35	18 - 73	6.2	-0.023	-0.027	1.8	suggestive		-
time spent											
moving	%	2	66	58 - 77	6	0.012	0.029	1.7	-		-
rearing	s	2	76	62 - 82	6	0.9	4	1.7	-		-
locomotion											
speed in											
light											
compartment											
nt	cm / s	7	24	7 - 52	5.4	2.2	-3.3	1.5	-		suggestive
locomotion											
speed in											
light											
compartment											
nt	cm / s	18	0	0 - 14	5.3	-0.3	2.5	1.5	-		-
fecal boli	count	18	21	2 - 25	8.3	0.5	0.3	2.4	suggestive		suggestive
fecal boli	count	4	58	52 - 71	6.5	0.5	-0.06	1.9	-		-
fecal boli	count	13	19	8 - 28	5	-0.3	-0.5	1.4	-		-

*Startle response*

startle											
response	g	2	47	26 - 57	7.6	30	-81	2.1	-		-
startle											
response	g	4	73	63 - 86	6	-32	-64	1.6	-		significant
startle											
response	g	8	26	17 - 38	5.7	43	-32	1.6	-		-
startle											
response	g	13	15	3 - 22	5.2	50	23	1.4	-		-

*Anatomy*

adrenal											
gland	mg	11	42	34 - 54	7.1	1.4	-1.9	1.8	suggestive		-
adrenal											
gland	mg	2	48	37 - 56	6.1	0.6	2.1	1.6	-		-

adrenal											
gland	mg	6	65	47 - 82	6.2	-1.4	1.3	1.6	-	-	
adrenal											
gland	mg	8	64	42 - 79	6.1	1.5	0.4	1.6	-	suggestive	
adrenal											
gland	mg	16	18	15 - 26	5.6	1.4	1.5	1.4	-	-	
adrenal											
gland	mg	12	11	6 - 25	5	-1.1	1.3	1.3	-	-	
body											
weight	g	18	16	0 - 22	6.8	6.3	-3.8	1.8	suggestive	-	
heart	g	18	20	6 - 26	5.4	0.03	-0.005	1.5	-	-	
kidney	g	3	54	31 - 66	6.9	0.02	0.01	2	-	-	
kidney	g	7	13	0 - 25	5.9	0.02	0.02	1.8	-	-	
kidney	g	8	88	85 - 97	5.9	0.02	0.01	1.7	-	-	
kidney	g	20	26	11 - 42	4.3	0.003	0.007	1.3	-	suggestive	
lung	g	10	53	35 - 67	5.8	0.02	-0.1	1.7	-	-	
lung	g	6	35	19 - 43	5.2	-0.007	0.06	1.5	-	-	
spleen	g	20	21	10 - 31	6.9	0.007	0.019	1.8	suggestive	-	
spleen	g	3	63	51 - 75	6.6	-0.015	0.014	1.8	-	-	
spleen	g	18	17	11 - 24	5.4	0.014	-0.005	1.5	-	suggestive	
testis	g	10	92	86 - 94	6.4	0.06	0.26	3.9	n/a	n/a	
testis	g	17	21	5 - 25	7.2	0.09	-0.09	4.3	n/a	n/a	
spotting	-	6	48	39 - 56	5.8	0.06	-0.14	1.8	-	suggestive	
<i>Serum traits</i>											
albumin	g / l	3	5	0 - 22	7.5	0.47	-0.66	2.3	significant	-	
albumin	g / l	4	62	52 - 85	5.4	-0.46	-0.17	1.6	-	-	
AP	µkat / l	10	52	8 - 72	5.8	-0.06	0.15	1.7	-	-	
AP	µkat / l	7	92	78 - 103	5.3	0.02	-0.11	1.6	-	-	
creatinine	µkat / l	15	23	15 - 28	6.9	-0.97	-0.67	2.1	-	-	
creatinine	µkat / l	19	51	30 - 62	5	-1.03	-1.10	1.5	-	-	
corticostero											
ne	ng / ml	5	18	1 - 30	6	-23.8	201	1.9	significant	-	
corticostero											
ne	ng / ml	6	88	64 - 94	6	-110	29	1.9	-	-	
fT4	pmol / l	5	94	76 - 109	8.5	-2.2	0.4	2.6	-	suggestive	
fT4	pmol / l	14	34	3 - 47	7.1	-1.8	-0.5	2.2	suggestive	-	
fT4	pmol / l	13	2	0 - 22	5.3	1.3	-1	1.6	-	-	
ALAT	µkat / l	5	67	57 - 77	7.3	-0.015	-0.045	2.3	-	-	
ALAT	µkat / l	12	21	10 - 42	5.8	0.026	-0.024	1.8	-	-	
ASAT	µkat / l	8	87	81 - 92	7.8	-0.14	0.25	2.4	suggestive	-	
glucose	µmol / l	2	49	37 - 57	6.6	0.67	-0.09	2	-	-	
urea	mmol / l	7	106	94 - 106	5.8	-0.06	-0.37	1.8	-	-	
total											
protein	g / l	3	117	112 - 17	6.3	0.75	0.38	2	-	significant	

total										
protein	g / l	11	15	7 - 19	6	0.28	-1.2	1.9	suggestive	-
total										
protein	g / l	7	64	50 - 103	5.8	0.6	0.65	1.8	-	-
total										
protein	g / l	8	2	0 - 36	5.6	-0.72	-0.86	1.7	-	-
total										
protein	g / l	6	89	66 - 94	5.2	-0.64	-0.56	1.6	-	-
Cl-	mmol / l	2	144	134 - 147	6.3	-0.7	0.67	1.9	suggestive	-
Cl-	mmol / l	10	46	10 - 71	5.4	-0.51	-0.27	1.7	-	-
Fe2+/3+	μmol / l	2	96	87 - 117	7	-2.4	-0.1	2.2	suggestive	-
Fe2+/3+	μmol / l	18	22	0 - 26	5.4	-0.02	1.8	1.7	-	-

Shown are QTLs with chromosome-wide (but not genome-wide) significance

**TABLE S6**  
**X-Chromosomal QTLs**

Trait	Peak (cM)	d.f. <sup>1</sup>	LR <sup>2</sup>	p value	additive effect	% variance <sup>3</sup>
urea	47	2	13.9	0.00096	-0.34	2.2

<sup>1</sup> degrees of freedom; <sup>2</sup> likelihood ratio; <sup>3</sup> residual phenotypic variance explained after accounting for fixed effects

**TABLE S7****Sex-specific QTLs identified at genome-wide significance**

Trait	sex	chr <sup>1</sup>	peak <sup>2</sup>	1 LOD c.i. <sup>2,3</sup>	F	additive effect	dominance	% variance <sup>4</sup>
adrenal gland	females	5	48	44 – 58	9.5	-2.7 mg	-1.3 mg	5.3
corticosterone	females	12	0	0 - 2	8.5	12 ng / ml	280 ng / ml	4.9

Only significant QTLs that do not overlap with QTLs found in the sex-combined analyses are shown. See Table 3 and Table S5 for sex-specific QTLs overlapping QTLs found in the sex-combined analyses. <sup>1</sup> chromosome; <sup>2</sup> cM; <sup>3</sup> confidence interval; <sup>4</sup> residual phenotypic variance explained after accounting for fixed effects;